

# Bioactivity of *Apis dorsata* Nest Extract from a Different Geographical Location in Minahasa, North Sulawesi, Indonesia

*by* Yermia Mocosuli

---

**Submission date:** 05-Jan-2023 07:04AM (UTC+0700)

**Submission ID:** 1988670123

**File name:** 71795\_01\_doi\_15.1997\_IJAB-22-0117\_8\_1-8\_1.pdf (309.13K)

**Word count:** 4716

**Character count:** 24466



## Full Length Article

# Bioactivity of *Apis dorsata* Nest Extract from a Different Geographical Location in Minahasa, North Sulawesi, Indonesia

Mokosuli Yernia Samuel<sup>1\*</sup> and Dina O. Rombot<sup>2</sup>

<sup>1</sup>Department of Biology, Faculty of Mathematics and Natural Science, State University of Manado, Tondano, Sulawesi Utara, Indonesia

<sup>2</sup>Faculty of Medicine, Sam Ratulangi University Manado, Sulawesi Utara, Indonesia

\*For correspondence: yermiamokosuli@unima.ac.id

Received 16 March 2022; Accepted 01 September 2022; Published 30 December 2022

## Abstract

The content of *Apis dorsata* Binghami nesting secondary metabolites differs based on the availability of plant feed sources. The present study was conducted to test for *in vivo* Brine Shrimp Lethality Test (BSLT) and radical scavenging activity of the ethanolic extracts of the nest. Nests were obtained from various natural habitats in the Minahasa area. Cytotoxicity was evaluated in LC<sub>50</sub> (lethality concentration 50), while radical scavenging activity was evaluated in IC<sub>50</sub> (inhibitory concentration). This research was conducted in 2021–2022. Results showed that the ethanolic extracts of a nest from North Minahasa had a higher phytochemical content intensity than other nest extracts from Minahasa. The DPPH free radical scavenging activity of all the ethanol extract of the nest from Minahasa was stronger than the positive control of ascorbic acid. The best IC<sub>50</sub> was shown by the nest ethanol extract from North Minahasa (57.67 mg/mL). The toxicity of nest ethanol extract was below 500 mg/mL, where the highest LC<sub>50</sub> was shown by North Minahasa nest extract (208.498 mg/mL). However, the extract toxicity based on the BSLT test was moderate. Thus, nest extract has the potential to be developed as a bioactive source of herbal medicine. © 2022 Friends Science Publishers

**Keywords:** Nest extract; Antioxidant; LC50; Cytotoxicity

## Introduction

A honey bee nest consists of a hexagonal-shaped chamber held together by Propolis and beeswax. There are eggs, larvae, pollen and royal jelly (Hadisoesoilo 2001). Propolis is one of the components that makes up the nest, which is widely studied today. This study used *Apis dorsata* Binghami nest, Sulawesi endemic honey bees (Hadisoesoilo 2001; Raffiudin 2002; Samuel *et al.* 2013). The nest of *A. dorsata* consists of only one comb. It is only found in the natural forests of Sulawesi and nearby small islands because it cannot be domesticated (Raffiudin 2002). As a giant honey bee, *A. dorsata* variety and types of food sources are greater than other honey bees (Repi and Samuel 2016; Samuel *et al.* 2013, 2019).

Propolis is a mixture of natural resins collected by bees from various plant species. Bee propolis is the product of plant resin processing with bee saliva, which contains various enzymes and peptides to produce new resins (Przybyłek and Karpinski 2019). Honey bees collected plant resins from plant parts, shoots and exudates (Wagh 2013; Kurek-Gorecka *et al.* 2014; Abdullah *et al.* 2020). Propolis also contains high micronutrients such as

vitamins (A, B, C), minerals (Ca, Cu, Fe, Mg, Mn, Na, Fe and Zn) and the enzyme succinate dehydrogenase (Salatino *et al.* 2011). The most secondary metabolites in Propolis are polyphenols (phenolic acids, flavonoids and their esters), steroids, terpenoids and amino acids (Sforcin and Bankova 2011; Kurek-Gorecka *et al.* 2014). Propolis has a colour and active content that varies depending on the plant source (Sforcin and Bankova 2011; Salatino *et al.* 2011; Abdullah *et al.* 2020). The natural resin substance in Propolis provides a sticky aroma and texture at the temperature of the newly formed nest. The health benefits of Propolis include accelerating cell regeneration, repairing leakage of blood vessels, curing diabetes mellitus, immune system stimulator, anticancer, anti prostaglandin, antistress, antioxidant, anti-inflammatory, antibacterial and antihyperlipidemic (Hegazi 1998; Bankova *et al.* 2000; Simone-Finstrom and Spivak 2010; Kubiliene *et al.* 2015). In addition, Propolis in the nest is very potential as an antiviral agent for SARS COV2 (Berretta *et al.* 2020).

Previous research has shown that *A. dorsata* nest extract from Minahasa can reduce the blood lipid profile of rats treated with hyperlipidemia (Mokosuli *et al.* 2019).

Analysis of flavonoid content using the HPLC method of Sarang *A. dorsata* extract obtained from North Minahasa in September obtained 23 types of flavonoid derivatives (Semuel *et al.* 2019). The bioactive content of *A. dorsata* nest highly depends on the type of feed available. Thus, it is necessary to research the bioactive content of *A. dorsata* nest from Minahasa obtained from various locations and different flowering seasons. The initial bioassay that can be done to determine the pharmacological potential of bioactive natural ingredients that is widely used is the Brine Shrimp Lethality Test (BSLT). BSLT is a method for determining the cytotoxic spectrum of natural extracts (Hamrun *et al.* 2020). Research on the analysis of phytochemical content, antioxidants, and BSLT nests from various regions and habitats in the Minahasa peninsula has been carried out.

## Materials and Methods

### Honey nest samples

The nests of *A. dorsata* obtained from four locations on the Minahasa peninsula, namely South Minahasa Regency, Southeast Minahasa Regency, Minahasa Regency and North Minahasa Regency (Fig. 1a, b). The nest was obtained directly from the tree where *A. dorsata* made the nest (Fig. 1c). The method of taking the nest using fumigation techniques is assisted by professional forest honey bee seekers. Nests collected at various locations have the same age. The age of the nest used had three months. The nest sample collection was carried out in September 2021.

### Manufacture of simplicia

Fresh samples of *A. dorsata* nest were cleaned and then washed in running water. The nest was then dried at 25°C for two days. Next, the nest was dried using an oven for 30 min at 50°C. At the end, the nest was finely chopped then blended until it becomes a powder. Nest powder is used for extraction.

### Extraction

The extraction was carried out using the maceration method (Harborne 1996; Mokusuli *et al.* 2019). The solvent and nest powder ratio is 1: 4 (w/v), namely 50 grams of macerated nest powder with 200 mL of 70% ethanol (chemical-pharmaceutical). Erlenmeyer contains solvent and nest powder, covered with aluminum foil, then placed in an orbital shaker for 2 × 24 h at a speed of 50 rpm and a temperature of 25°C. The macerate is then filtered using Whatman filter paper to obtain the filtrate. Finally, the filtrate obtained is dissolved using a Buchi rotary evaporator at a speed of 55 rpm and a temperature of 50°C. Furthermore, the evaporation results are called the crude extract of the nest of *A. dorsata*.

### Phytochemical content analysis

Phytochemical content analysis using the Harborne (1996) method with slight modifications was done. The phytochemicals analyzed were alkaloids, flavonoids, saponins, tannins, steroids and triterpenoids.

### DPPH method free radical reduction activity test

*A. dorsata* nest extract was made to distribute concentrations (10, 50, 100, 200 and 250 µg/µL). Each one is put into a test tube. Each test tube added 500 µL of DPPH 1 mM solution in methanol. The volume was crushed to 5.0 mL, then incubated at 37°C for 30 min. Using a UV spectrophotometer (Parkin Elmer), the absorbance was measured at 515 nm. As a positive control, ascorbic acid (Sigma Aldric) was used with an adjusted concentration (Imazawa *et al.* 2004; Semuel 2008). The IC<sub>50</sub> values are calculated using the linear regression equation formula:

$$\% \text{ inhibition} = \frac{[\text{Absorbance control} - \text{absorbance of the sample}]}{\text{Absorbance control}} \times 100\%$$

### BSLT method cytotoxic test

The *A. salina* Leach cyst was weighed as much as 20 mg then put into a particular container containing filtered seawater; after being aerated, the cyst was left for 48 h under lamp lighting to hatch completely. The hatched larvae are taken for use in toxicity tests—a total of 10 healthy *A. Salina* Leach larvae (based on the motility and ability of the larvae to search for light) were put into the test vial containing seawater. The ethanol extract solution was added to each test vial with the concentration of the test solution consisting of 10, 100, 500, and 1000 ppm, while the control extract was not added to the extract solution. Each of the three replications was made. Observations were made after 24 h by counting the number of larvae that died from the total larvae included in the test vial (McLaughlin *et al.* 1998).


### Statistical Analysis

The calculation uses a magnifying glass. The LC<sub>50</sub> value was determined by probit analysis, with a 95% degree of significance using the IBM SPSS 20.

### Results

The nests of *A. dorsata* from Sawangan Minahasa Utara were golden yellow in color. *A. dorsata* was nesting in the stalk of the *Ficus* spp. The nest has a fruity aroma, especially Langsat (*Lansium domesticum* L.) fruit. Nest 1.3 meters wide, hanging from twigs 22 m from the ground. The honey taste still found in the nest is sweet with a slightly sour taste. From observations, around nesting in September, the flowering season of Langsat, durian, mango,






**Fig. 1a:** Locations for sampling *A. dorsata* nests are marked with an asterisk



**Fig. 1b:** Location of *A. dorsata* nest in Minahasa, North Sulawesi, Indonesia



**Fig. 1c:** Sample nest of *A. dorsata* from (a). Minahasa Utara (b). Minahasa Induk (c). Southeast Minahasa (d). Minahasa Selatan




4  
**Fig. 2:** Nest extract (a), Nest powder (b), Macerated filtrate (c). Solvent evaporation with a rotary evaporator at 450°C, 55 rpm (d). Crude extract

and other fruit plants. The nests of *A. dorsata* from Kombi Village, Minahasa, are brownish-yellow in colour. The age of the nest is about three months. The taste of honey found in the hive is sweeter than the honey in the North Minahasa nest. However, the hive and honey smelled like cloves. Nests were also found in the *Ficus* spp. tree, with a nest height of 19 meters from the ground. About 300 meters from the nest, there is a clove plantation where the cloves are in the stage of producing flowers and fruit in September. The nests of *A. dorsata* from Ratahan Village, Southeast Minahasa Regency, are brownish-yellow but more yellow than the main minahasa nests. Honey and fruit-scented nest. Nests were found in Aren trees, with 15 meters from the ground. Around the nest is still a natural forest with lots of Aren trees. The nest of *A. dorsata* from Wulurmaatus Village, Modinging Minahasa Selatan District, was found on a pedu tree (local name), at an altitude of 17 meters from the ground. The nests are dark brown, have a very sweet honey taste and have a fruity aroma (Fig. 2).

#### *A. dorsata* honey nest extract

Nest after mashed brownish yellow, scented with cloves. Nest mashed using a blender has been cleaned of larvae. The smell of cloves is caused by the nest location close to a clove plantation, around 700–1000 m. The filtrate resulting from nest maceration with 70% ethanol for 2 × 24 h has a brownish yellow colour (Fig. 2).



4  
**Fig. 3:** Free radical scavenging activity of nest extract

After the solvent was evaporated with the Buchi Rotary Evaporator, crude extract of *A. dorsata* nest from South Minahasa was brownish-black with a weight of 50 g. The crude extract had a distinctive nest aroma, honey, cloves and fruit aroma (Fig. 3).

#### Phytochemical content

Based on the analysis of phytochemical content, all nest samples were identified to contain all phytochemical groups. Nest ethanol extract from all locations in Minahasa contains high intensity flavonoids, saponins and tannins.

**Table 1:** Percentage of the yield of *A. dorsata* nest extract from various districts in North Sulawesi

Sample	Sample weight (g)	Extract weight (g)	Yield (%)
SMS	50	3.26	6.51
SMI	50	3.21	6.42
SMU	50	3.10	6.20
SMT	50	3.08	6.16

Description:

SMS = nest from South Minahasa : Modounding

SMI = nest from Minahasa : Kombi

SMU = nest from north Minahasa : Airmadidi

SMT = nest from southeast Minahasa : Ratahan

**Table 2:** Phytochemical content of the ethanol extract of *A. dorsata* nest from various regions in North Sulawesi

Phytochemical Group	SMS	SMI	SMU	SMT
Alkaloids	+	+	+	+
Flavonoids	++	++	++	++
Saponins	+++	++	+++	++
Tannins	++	++	++	++
Steroids				
- Meyer reagent	+	+	+	+/-
- Wagner reagent	+	+/-	+	+
- Dragendorff reagent	+	++	+/-	+
Triterpenoids				
- Meyer reagent	+	+/-	+	+/-
- Wagner reagent	+/-	+	+/-	+
- Dragendorff reagent	+	+	+	+

Description:

+: there are groups of phytochemical compounds based on the intensity of appearance after being given a reagent.

-: not available based on the intensity of appearance after being given reagent.

**Table 3:** Percentage of nest extract inhibition

Samples	Test concentration (mg/mL)				
	10	50	100	200	250
SMU	16.65	30.22	52.28	58.07	68.06
SMI	11.93	39.5	44.31	52.89	67.23
SMT	11.56	39.65	47.91	59.74	67.4
SMU	11.56	39.65	47.91	59.74	67.4
VIT C (control)	7.67	29.23	40.51	49.76	58.07

However, the content of steroids and triterpenoids is of little intensity. All nest extract samples identified contained alkaloids (Table 1–3).

#### Antioxidant test

The percentage of inhibition of the extract against DPPH free radicals varied based on the origin of the nest in Minahasa. The highest mean percentage of free radical reduction of all samples at the test concentration of 250 mg/mL. However, there was no significant difference in *A. dorsata* nest extract concentration from various districts in Minahasa. However, the inhibition percentage of extracts at the test concentration of 250 mg/L was higher than the control used, namely vitamin C (Table 3).

The highest IC<sub>50</sub> of the extract against DPPH free radicals was shown by the nest extract from North Minahasa (SMU), namely 57.67 mg/mL (R = 0.94). Furthermore, SMT nest extract is 100.83 mg/mL (R = 0.99), SMU nest

extract is 103.98 mg/mL (R = 0.94) and SMI nest extract is 118.2 mg/mL (R = 0.95). Compared with the control vitamin C which had an IC<sub>50</sub> of 179.06 mg/mL (R = 0.98), the IC<sub>50</sub> of nest extract from all locations in Minahasa was still better (Fig. 3).

#### Bioassay Toxicity Brine Shrimp Lethality Test (BSLT) method

The highest average mortality in all nest samples was shown in the extract treatment with a concentration of 1000 ppm. In contrast, the lowest average mortality was shown in the 10 ppm extract concentration treatment in all nest extract samples. The percentage of mortality of *Artemia salina* Leach larvae increases with increasing test concentration (Fig. 4). From the probit analysis, the graph of the extract mortality plots is in linear regression (Fig. 5).

Based on the probit analysis of the mortality of *Artemia salina* Leach after application of nest extract, the highest LC<sub>50</sub> was shown by SMU extract (208.498 mg/mL). In comparison, the extract with the lowest LC<sub>50</sub> was SMS extract (395.186 mg/mL). Based on the LC<sub>50</sub> values obtained by the SMS, SMI and SMU extracts were less than 500 mg/mL (Fig. 6).

#### Discussion

The *A. dorsata* nests in Minahasa have different colours, honey taste and aroma characteristics. However, the average length of the nest was relatively the same at the age of 2–3 months, namely 1–1.75 meters and is semicircular (Semuel *et al.* 2019) The differences in taste, aroma and colour are influenced by the forage plants, which are the source of nectar, pollen, plant resins around the nest. The *A. dorsata* bee has a greater variety of forage plants than *A. mellifera*, with a flight distance of about 600–900 meters from the nest (Semuel *et al.* 2013). Nests from North Minahasa are in areas adjacent to plantations and mountainous areas, while nests from Minahasa are in areas adjacent to the sea. Compared to the nests from South Minahasa, which are in the highlands with cooler temperatures and nests from Southeast Minahasa, which are found in mountainous areas and far from people's plantations. Thus, the characteristics of *A. dorsata* hives and honey are also influenced by geographic location (Semuel *et al.* 2013; Semuel and Repi 2015; Semuel *et al.* 2019).

The characteristics of the nest also positively correlate with the characteristics of the phytochemical content. The results of the nest phytochemical group analysis found variations in the intensity of the content of each phytochemical class of *A. dorsata* hives from various regions in Minahasa. The highest percentage of DPPH and IC<sub>50</sub> free radical inhibition was the nest samples from North Minahasa. In contrast, the DPPH and IC<sub>50</sub> free radical inhibition of the nest samples from Southeast Minahasa and South Minahasa were not significantly different. Based on




Fig. 4: Comparison of the percentage of mortality of *Artemia salina* Leach larvae based on the origin of the nest sample




Fig. 5: Graph of LC<sub>50</sub> probit analysis of *A. dorsata* nest extract




Fig. 6: LC<sub>50</sub> of nest extract based on sample origin from several districts in Minahasa



the analysis of the phytochemical content of the *A. dorsata* nests from North Minahasa, had a higher intensity than the nest samples from other areas in Minahasa (Semuel *et al.* 2019). However, the IC<sub>50</sub> of the nest extract was below 150 ppm or better than the positive control for ascorbic acid. Thus, the nest extract's DPPH free radical scavenging activity was different based on the origin of the nest samples in Minahasa.

Based on the results of the BSLT test, the toxicity of the nest extracts of SMS, SMI, SMT and SMU were below 500 mg/mL, thus the toxicity of the nest extract was in the moderate category in *A. salina* Leach. The high-intensity content of phytochemicals, namely flavonoids, saponins and tannins, affects the toxicity of the extract. The active compound in nest propolis is soluble in ethanol (Kubiliene *et al.* 2015). These phenolic group compounds are cytotoxic in *A. salina* Leach larvae (Semuel 2008; Ullah *et al.* 2013; Marzuki *et al.* 2019; Rasyid *et al.* 2020). The active compound activity in the extract that enters the larvae and goes to the cells, causing metabolic and cytotoxic disorders. Cytotoxic effects can be observed quickly within 24 h (McLaughlin 1998; Dwijayanti *et al.* 2015; Rachman *et al.* 2020). Phenolic compounds can induce apoptosis in cells (Semuel 2008). Tunisian ethanolic extract of Propolis exhibits strong antioxidant, antibacterial and antiproliferative activity (Béji-Srairi *et al.* 2020). BSLT has been used by many researchers, including initial toxicity testing, including fungal extracts (Carballo *et al.* 2002), langsat stem bark extract (Semuel 2008), Averhoa bilimbi leaf extract (Rachman *et al.* 2020), metal weight and pesticides (Hussain *et al.* 2006), *Arecha catechu* seed extract (Rasyid *et al.* 2020), *Ruellia tuberosa* L. leaf extract (Vitalia *et al.* 2016) and rosella flower extract (Purbowati *et al.* 2015). Shrimp larvae are known to have thin skin to be sensitive to their environment. Foreign substances or compounds that exist in the environment will diffuse into the body so that they affect the cells. If the compound that enters the cell is toxic, it will kill the shrimp larvae. According to Carballo *et al.* (2002), an extract of natural ingredients has potential as medicine if it has an LC<sub>50</sub> according to the BSLT test of less than 1000 mg/mL. The toxicity of natural extracts based on LC<sub>50</sub> values is very strong (< 10 ppm), strong (10–100 ppm), moderate (100–500 ppm) and weak (500–1000 ppm) (Meyer *et al.* 1982; McLaughlin *et al.* 1998).

The propolis extract from Bangladesh honey bees obtained LC<sub>50</sub> 57.99 (Tanvir *et al.* 2018), compared with the crude extract of honey bee hive in this study, LC<sub>50</sub> honey nest extract of *A. dorsata* has the potential to be developed as a bioactive source that has pharmacological properties. The nest extract of *A. cerana* showed strong toxicity (LC<sub>50</sub> 67,744 and 86,98 (Yasmin *et al.* 2019); Propolis from *Trigona itama* from Beladin Serawak showed low toxicity but high free radical scavenging activity (EC<sub>50</sub> 17.18 mg/mL) (Yusop *et al.* 2019). In previous research, the nest ethanol extract from North Minahasa showed strong antioxidant activity with IC<sub>50</sub> 6.69 mg/mL, while n-hexane

extract had IC<sub>50</sub> 6.76 mg/mL (Semuel *et al.* 2019). The IC<sub>50</sub> obtained was greater in this study but still had a strong free radical scavenging activity > 200 mg/mL. Season and diversity of bee food sources (Semuel *et al.* 2013, 2019).

Empirically, the Minahasa people consume bee hives directly because they are believed to have medicinal properties. Bee hives are believed to reduce blood cholesterol, uric acid, treat diabetes and have anti-tumor properties (Kaunang and Mocosuli 2017). The medium toxicity category indicates that the active compounds present in the nest extract from all locations in Minahasa have pharmacological activities. With the LC<sub>50</sub> in the moderate category it confirms that nest extract can be consumed directly by humans.

## Conclusion

The analysis of the toxicity test of *A. dorsata* nest extract had LC<sub>50</sub> 54.457. Based on the LC<sub>50</sub> value, the *A. dorsata* forest honey nest can be consumed directly by humans due to containing bioactive medicinal potential.

## Acknowledgement

Thanks to Directorate of Research and Community Service, Ministry of Research, Technology and National Innovation Research Agency of the Republic of Indonesia for funding this research through the PDUPT scheme in 2021.

## Author Contributions

Author Contributions MYA designed the experiment, collected and prepared the materials, analyzed the data and drafted the manuscript. DR prepared the nest extracts, collected data and reviewed the manuscript. MYA and DR collected data and revised the manuscript.

## Conflicts of Interest

The authors declare no conflict of interest.

## Data Availability

Data presented in this study will be available on a fair request to the corresponding author.

## Ethics Approval

Not applicable in this paper.

## Reference

Abdullah NA, N Zulkiflee, SNZ Zaini, H Taha, F Hashim, A Usman (2020). Phytochemicals, mineral contents, antioxidants, and antimicrobial activities of Propolis produced by Brunei stingless bees *Geniotrigona thoracica*, *Heterotrigona itama* and *Tetrigona binghami*. *Saud J Biol Sci* 27:2902–2911



- Bankova VS, SLD Castrob, MC Marcucci (2000). Propolis: Recent advances in chemistry and plant origin. *Apidologie* 31:3–15
- Béji-Srairi R, I Younes, M Snoussi, K Yahyaoui, G Borchard, R Ksouri, MK Wided (2020). Ethanol extract of Tunisian Propolis: Chemical composition, antioxidant, antimicrobial and antiproliferative properties. *J Apicul Res* 59:917–927
- Berretta AA, MAD Silveira, JMC Capcha, DD Jong (2020). Propolis and its potential against SARS-CoV-2 infection mechanisms and COVID-19 disease. *Biomed Pharmacother* 131:110622
- Carballo JL, ZL Hernandez-Inda, P Perez, MD Garcia-Gravalos (2002). A comparison between two brine shrimp assays to detect *in vitro* cytotoxicity in marine natural products. *BMC Biotechnol* 2:1–5
- Dwijayanti E, HA Andi, MA Wibowo (2015). Skrining fitokimia dan uji aktivitas sitotoksik pada kulit batang tampoi (*Baccaurea macrocarpa*) terhadap *Artemia salina* Leach dengan Metode BSLT. *J Kim Khatul* 4:6–10
- Hadiosesilo S (2001). Keanekaragaman spesies lebah madu asli Indonesia. *Biodiversitas* 2:123–128
- Hamrun N, T Nabilah, R Hasyim, M Ruslin, I Dammar, MA As (2020). Toxicity test of bioactive red alga extract *Eucheuma spinosum* on shrimp *artemia salina* leach. *Syst Rev Pharm* 11:672–676
- Harborne JB (1996). *Metode Fitokimia Penuntun Cara Modern Menganalisis Tumbuhan*. ITB, Bandung, Indonesia
- Hegazi AG (1998). Propolis an overview. *J Bee Inform* 5:22–28
- Hussain A, M Zia, B Mirza (2006). Cytotoxic and antitumor potential of *Fagonia cretica* L. *Turk J Biol* 31:19–24
- Kaunang ESN, YS Mokusuli (2017). Botanical and phytochemical constituents of several medicinal plants from mount Klabat north Minahasa. *J Med Plants Stud* 5:29–35
- Kubiliene L, V Laugaliene, A Pavilonis, A Maruska, D Majiene, K Barcauskaite, A Savickas (2015). Alternative preparation of propolis extracts: Comparison of their composition and biological activities. *BMC Complem Altern Med* 15:156–162
- Kumazawa S, T Hamasaka, T Nakayama (2004). Antioxidant activity of Propolis of various geographic origin. *Food Chem* 84:329–339
- Kurek-Gorecka A, A Rzepecka-Stojko, M Górecki, J Stojko, M Sosada, G Świerczek-Zięba (2014). Structure and antioxidant activity of polyphenols derived from Propolis. *Molecules* 19:78–101
- Marzuki A, L Rahman, SS Mamada (2019). Toxicity test of stem bark extract of banyuru (*Pterospermum celebicum* Miq.) using BSLT (brine shrimp lethality test) and cream irritation test. *J Phys Conf Ser* 1341:1–9
- McLaughlin JL, LL Rogers, JE Anderson (1998). The use of biological assays to evaluate botanicals. *Drug Inform J* 32:513–524
- Meyer BN, NR Ferrigni, JE Putnam, LB Jacobsen, DEJ, Nichols, JL McLaughlin (1982). Brine shrimp: A convenient general bioassay for active plant constituents. *Plant Med* 45:31–34
- Przybyłek I, TM Karpinski (2019). Antibacterial properties of propolis. *Molecules* 24:2047–2063
- Purbowati ISM, K Syamsu, E Warsiki, HS Rukmini (2015). Evaluasi toksisitas, aktivitas antibakteri dan antioksidan komponen bioaktif rosela dengan variasi jenis pelarut. *J Agroindustr Technol* 25:1–7
- Rachman SR, F Maryam, BA Limbong (2020). The toxicity test of ethanol ekstrak of leaves *Averhoa bilimbi* L. Using Brine Shrimp Lethality Test (BSLT) Methode on shrimp larvae (*Artemia salina* Leach). *J Pharm Med Sci* 4:1–5
- Raffiudin R (2002). Honey bee behavioural evolution and itr gene structure studies. *Doctoral dissertation*. James Cook University, Townsville, Queensland, Australia
- Rasyid MI, H Yuliani, L Angraeni (2020). Toxicity test LC50 of pineung nyen teusalee seeds (*Areca catechu*) Extract by Brine Shrimp Lethality Test (BSLT) Methode. *IOP Conf Ser: Earth Environ Sci* 515:12052
- Salatino A, CC Fernandes-Silva, AA Righi, MLF Salatino (2011). Propolis research and the chemistry of plant products. *Nat Prod Rep* 28:925–936
- Semuel MY (2008). Aktivitas antioksidan dan antikanker ekstrak kulit buah batang langsung (*Lansium Domesticum* L.). Thesis. Sekolah Pasca Sarjana Institut Pertanian Bogor, Bogor Regency, Java, Indonesia
- Semuel MY, Repi RA (2016). The Potential of herbal plant extracts native to South Minahasa as traditional coconut oil anti rancid Agents. *J Agric Health* 6:77–85
- Semuel MY, RA Repi (2015). The characteristics of bioactive peptides and antibacterial activity of honey bee (*Apis nigrocincta*) Smith Venom, endemic to Sulawesi. *Molekul* 10:135–144
- Semuel MY ESN Kaunang, JS Manopo (2019). The bioactive contents and antioxidant activity of honey bee nest extract of *Apis dorsata* Binghami from the North Sulawesi. *Molekul* 14:92–102
- Semuel MY, J Pelealu, M Tulung, LC Mandey (2013). Pharmacological bioactivity honey bee venom *Apis nigrocincta* Smith and *A. dorsata* Binghami endemic to North Sulawesi. *Intl J Sci Eng Invest* 2:25–33
- Seorcin JM, V Bankova (2011). Propolis: Is there a potential for the development of new drugs?. *J Ethnopharmacol* 133:253–260
- Simone-Finstrom M, M Spivak (2010). Propolis and bee health: The natural history and significance of resin use by honey bees. *Apidologie* 41:295–311
- Tanvir EM, MS Hossen, UM Shapla, M Mondal, R Afroz, M Mandal, SH Gan (2018). Antioxidant, brine shrimp lethality and analgesic properties of Propolis from Bangladesh. *J Food Biochem* 42:1–12
- Ullah MO, M Haque, KF Urmi, AHM Zulfiker, ES Anita, M Begum, K Hamid (2013). Antibacterial activity and brine shrimp lethality bioassay of methanolic extracts of fourteen different edible vegetables from Bangladesh. *Asian Pac J Trop Biomed* 3:1–7
- Vitalia N, A Najib, AR Ahmad (2016). Uji toksisitas ekstrak daun pletekan (*Ruellia tuberosa* L.) dengan menggunakan metode Brine Shrimp Lethality Test (BSLT). *J Fitofar Indo* 3:124–129
- Wagh VD (2013). Propolis: A wonder bees product and its pharmacological potentials. *Adv Pharmacol Sci* 2013:1–11
- Yasmin Y, L Fitri, Fauziah, F Waslyyah (2019). Isolation of bacteria from *Apis cerana* hive, their antibacterial potency and cytotoxicity. *Biodivers J Biol Divers* 20:2733–2738
- Yusop SATW, AH Sukairi, WMAW Sabri, MR Asaruddin (2019). Antioxidant, antimicrobial and cytotoxicity activities of Propolis from Beladin, Sarawak Stingless Bees *Trigona itama* Extract. *Mater Today Proc* 19:1752–1760

# Bioactivity of Apis dorsata Nest Extract from a Different Geographical Location in Minahasa, North Sulawesi, Indonesia

## ORIGINALITY REPORT

14%

SIMILARITY INDEX

12%

INTERNET SOURCES

7%

PUBLICATIONS

%

STUDENT PAPERS

## PRIMARY SOURCES

1	<a href="http://www.fspublishers.org">www.fspublishers.org</a> Internet Source	5%
2	Nasriadi Dali Wela, Seniwati Dali, Armadi Chairunnas, Hilda Ayu Melvi Amalia, Sri Ayu Andini Puspitasari. "Extraction of The Chemical Components of Dengen Leaves ( <i>Dillenia serrata</i> Thunb) by MAE Method and Activity Test as Antioxidant and Toxicity", <i>Indo. J. Chem. Res.</i> , 2022 Publication	1%
3	<a href="http://smujo.id">smujo.id</a> Internet Source	1%
4	<a href="http://ojs.jmolekul.com">ojs.jmolekul.com</a> Internet Source	1%
5	<a href="http://repository.unima.ac.id:8080">repository.unima.ac.id:8080</a> Internet Source	1%
6	Wellem Anselmus Teniwut, Syahibul Kahfi Hamid, Marvin Mario Makailipessy. "Developing a masterplan for a sustainable	1%

marine sector in a small islands region:  
Integrated MCE spatial analysis for decision  
making", Land Use Policy, 2022

Publication

---

7	<a href="http://ijper.org">ijper.org</a> Internet Source	1 %
8	<a href="http://fmipa.unsrat.ac.id">fmipa.unsrat.ac.id</a> Internet Source	1 %
9	<a href="http://ejurnal-mapalus-unima.ac.id">ejurnal-mapalus-unima.ac.id</a> Internet Source	<1 %
10	<a href="http://ejournal.undiksha.ac.id">ejournal.undiksha.ac.id</a> Internet Source	<1 %
11	Biva, Israt Jahan, Md Mynol Islam Vhuiyan, Moni Rani Saha, Muhammad Shahidul Islam, and Shammy Sarwar. "In-vitro Antioxidant and Cytotoxicity Studies of <i>Annona squamosa</i> Linn.", Stamford Journal of Pharmaceutical Sciences, 2010. Publication	<1 %
12	Reymon Reymon, Suhikma Sofyan, Agung Wibawa Mahatva Yodha, Musdalipah Musdalipah. "The toxicity of <i>Meistera chinensis</i> rhizome fraction by shrimp larvae with the BSLT method", Natural Science: Journal of Science and Technology, 2021 Publication	<1 %

---



13 Pooran Golkar, Fariborz Moattar. " Essential Oil Composition, Bioactive Compounds, and Antioxidant Activities in L. ", Natural Product Communications, 2019 <1 %  
Publication

---

14 innovareacademics.in <1 %  
Internet Source

---

15 res.mdpi.com <1 %  
Internet Source

---

16 www.spandidos-publications.com <1 %  
Internet Source

---

17 strathprints.strath.ac.uk <1 %  
Internet Source

---

18 investor-id.jasamarga.com <1 %  
Internet Source

---

19 journalofnaturalproducts.com <1 %  
Internet Source

---

20 openjournals.ljmu.ac.uk <1 %  
Internet Source

---

Exclude quotes On

Exclude matches < 10 words

Exclude bibliography On