

ISBN. 978-979-99314-8-1

INTERNATIONAL CONFERENCE ON  
RESEARCH, IMPLEMENTATION AND  
EDUCATION OF MATHEMATICS AND  
SCIENCES 2014



# ICRIEMS 2014

Yogyakarta, 18-20 May 2014

*Global Trends and Issues  
on Mathematics and Sciences  
and the Education*

**PROCEEDING**

11  
15

Conference Proceedings

---

INTERNATIONAL CONFERENCE ON RESEARCH,  
IMPLEMENTATION AND EDUCATION  
OF MATHEMATICS AND SCIENCES (ICRIEMS) 2014  
Yogyakarta, 18 – 20 May 2014

---

ISBN 978-979-99314-8-1



Global Trends and Issues  
on Mathematics and Science  
and The Education

Faculty of Mathematics and Natural Sciences  
Yogyakarta State University

Conference Proceedings  
ICRIEMS 2014 : Global Trends and Issues on Mathematics and Science  
and The Education

- Mathematics & Mathematics Education
- Physics & Physics Education
- Chemistry & Chemistry Education
- Biology & Biology Education
- Science Education

© June 2014

**Editorial Board:**

Hari Sutrisno  
Wipar Sunu Brams Dwandaru  
Kus Prihantoso Krisnawan  
Denny Darmawan  
Erfan Priyambodo  
Evy Yulianti  
Sabar Nurohman

**Board of Reviewer**

Prof. Dr. Saberi bin Othman (Universiti Pendidikan Sultan Idris, Malaysia)  
Prof. Samsuuk Ekasit (Mahidol University, Thailand)  
Prof. Dean Zollman (Kansas State University, U.S.)  
Prof. Tran Vui (Hue University, Vietnam)  
Prof. Dr. Amy Cutter-Mackenzie (Southern Cross University, Australia)  
Dr. Alexandra Lynman (Universitas Hindu Indonesia)  
Prof. Dr. Rusgianto Heri Santoso (Yogyakarta State University)  
Prof. Dr. Marsigit (Yogyakarta State University)  
Prof. Dr. Mundilarto (Yogyakarta State University)  
Prof. Dr. Zuhdan Kun Prasetyo (Yogyakarta State University)  
Prof. Dr. K.H. Sugijarto (Yogyakarta State University)  
Prof. Dr. A.K. Prodjosantoso (Yogyakarta State University)  
Prof. Dr. Djukri (Yogyakarta State University)  
Prof. Dr. Bambang Subali (Yogyakarta State University)



Research  
may be  
the artic  
already  
Mathem  
of 344 p  
physics  
and scie

Science  
speaker  
Prof. D  
Macker  
(Hue U  
Mahid  
Tenaga  
12 uni  
speake  
Beside  
the lat  
These  
includ

studies  
access

## **Preface**

Bless upon God Almighty such that this proceeding on International Conference on Research, Implementation, and Education of Mathematics and Sciences (ICRIEMS) 2014 may be compiled according to the schedule provided by the organizing committee. All of the articles in this proceeding are obtained by selection process by the reviewer team and already been presented in the Conference on 18 – 20 May 2014 in the Faculty of Mathematics and Natural Sciences, Yogyakarta State University. This proceeding consists of 344 parallel papers, and comprises 9 fields, that is mathematics, mathematics education, physics, physics education, chemistry, chemistry education, biology, biology education, and science education.

The theme of ICRIEMS 2014 is 'Global Trends and Issues of Mathematics and Science and the Education'. The main articles in this conference are given by five keynote speakers, which are Prof. Dean Zollman (Physics Department, Kansas State University), Prof. David F. Treagust (Center of Education, Curtin University), Prof. Dr. Amy Cutter-Mackenzie (School of Education, Southern Cross University, Australia), Prof. Tran Vui (Hue University, Vietnam), and Asst. Prof. Dr. Duangjai Nacapricha (Faculty of Science, Mahidol University). The conference is also supported by the LPTK (Lembaga Pendidikan Tenaga Kependidikan) Forum from Faculty of Mathematics and Sciences that consists of 12 universities all over Indonesia. Each member of the Forum contributed one invited speakers, such that there are an additional 10 invited speakers presenting in the forum. Besides the keynote and invited speakers, there are also 344 parallel articles that presented the latest research results in the field of mathematics and sciences, and the education. These parallel session speakers come from researchers from Indonesia and abroad, including Malaysia and Australia.

Hopefully, this proceeding may contribute in disseminating research results and studies in the field of Mathematics and Sciences and the Education such that they are accessible by many people and useful for the Nation Building.

Yogyakarta, June 2014

The Editor Team

### Forewords from The Head of Committee

*Assalamu'alaikum wa Rahmatullahi wa Barakatuh*  
May God bless upon us.

Your excellency The president of UNY Prof. Dr. Rochmat Wahab, M. Pd., M.A., ladies and gentlemen, good morning and welcome to State University Yogyakarta. This seminar entitled International Conference on Research, Implementation, and Education of Mathematics and Science (ICRIEMS): global trends and issues on mathematics and science and the education is organized by the Faculty of Mathematics and Science, State University of Yogyakarta working together with 12 members of the Association of the Faculty of Math and Sciences from Teacher Education Program (LPTK). This seminar is also dedicated to the golden anniversary of UNY; 1 among 90 academic activities dedicated to the anniversary.



Ladies and gentlemen, on behalf of the committee of this conference, I would like to express highest appreciation and gratitudes to the keynote speakers, including:

1. Prof. David F. Treagust (Center of Science Education Curtin University)
2. Prof. Dean Zollman (Physics Dept, Kansas University, US)
3. Dr. Amy Cutter-Mackenzie (School of Education, Southern Cross University, Australia)
4. Asst. Prof. Dr. Duangjai Nacapricha (Faculty of Science , Mahidol University)
5. Prof. Tran Vui (College of Education, Hue University, Hue City, Vietnam)

Secondly, I would like also to give sincere thanks and gratitudes to the speakers from 10 College of Educations, including:

1. Universitas Negeri Surabaya (UNESA): Prof. Dr. Muchlas Samani, and 33 speakers
2. Universitas Negeri Jakarta (UNJ): Prof. Dr. Gerardus Pola, and 7 speaker
3. Universitas Pendidikan Indonesia (UPI): Dr. Hary Firman, and
4. Universitas Negeri Malang (UM): Prof. Effendi, Ph.D
5. Universitas Negeri Padang (UNP): Prof. Tjeerd Plomp
6. Universitas Negeri Semarang (UNNES): Prof. Dr. Supriyadi Rustad

7. Universitas Pendidikan Singaraja (UNDIKSA): Prof. Dr. I Nengah Suparta, M.Si
8. Universitas Negeri Makasar (UNM): Oslan Junaidi, Ph.D
9. Universitas Negeri Gorontalo (UNG): Prof. Dr. Sarson Pomalto, M.Pd
10. Universitas Negeri Yogyakarta (UNY): Dr. Jaslin Ikhsan

Next, I also would like to thanks to our special guests and speakers from:

1. Universitas Pendidikan Sultan Indris (UPSI), Malaysia
2. University of Mahidol, Thailand
3. University of Malaysia in Trengganu

Next, I would like to thanks and welcome to 379 speakers from the entire Indonesia and all participants registered in this seminar.

Ladies and gentlemen, recently the number of research and publication on mathematics and science and the education is vulnarable. It is nescessary for us to organise, to share, and to publish the results of the research in this conference. I hope the conference will bear fruitful results and promote networking and future collaborations for all participants from diverse background of expertise, intitutions, and countries to promote science, mathematics, and the education.

Finally, I am delighted to thank the committee members who have been working very hard to ensure the succes of the conference.

Please enjoy the conference and enjoy Yogyakarta, the city of education, tourism, and culture. Thank you very much.

Assalamu'alaikum wa rahmatullahi wa barrakatuh

**Dr. Slamet Suyanto, M. Ed.**

**STUDY ON POPULATION OF FROM *D. MELANOGASTER* KATUL MEDIA  
FERMENTED *SACHAROMYCES CEREVISIAE* AS SWIFLET WOOF**

**Ignatius R. S. Santoso**

<sup>1</sup>*Department of Chemistry, Faculty of Mathematics and Natural Sciences, Manado State  
University*

**ABSTRACT**

An identification of *D.melanogaster* population has been developed through artificial medium of fermented whole grain waste product by *Sacharomyces cereviceae*; which it's application and simpel technology is highly economic valuable for swift feed to enhance its quality and quantity. Fermented whole grain waste product has not been used as insect feed. In order to unravel the decreasing of fruits production during dry season which implicated the decreasing of *D.melanogaster* population that use as swift feed. This research aim to identify *D.melanogaster* population from banana, compare to artificial medium. Followed by chemical analysis of protein, carbohydrate, fat, mineral, and calorie value; determine the feed availability with swift nest population as well as determine the nutrition chemical compound in swift nest; mass, thickness, protein, carbohydrate, fat and mineral. The fact that fermented whole grain waste product can growth the *D.melanogaster* which is used for swift feed, swift farmers can develop *D.melanogaster* to increase quality and quantity of swift nest. Based on analysis of variance, natural medium from banana on population of *D.melanogaster* growth. For artificial feed, fermented whole grain waste product, the E treatment of *D.melanogaster* with banana fruit show the best result;

Keywords : *D.melanogaster*; swiftlet nest.

**A. INTRODUCTION**

The development of agriculture is not free from problems breathing pest and disease control, this damage occurs either on the ground, a warehouse, a lot of fruits and vegetables sold in traditional markets and supermarkets look clean and not deformed, but not the least the buyer fruit complain with poor quality fruit they buy. This fact is very influential in vegetable and fruit traders, because fruits and vegetables tersebut rapid deterioration and rotting (Novizan, 2002). One cause damage to fruit crops is the Fruit Fly *Drosophila* sp which is damaging the flesh of the fruit flies that memyebabkan ripe fruit becomes rotten and falling. Fruit flies can be the same size larger or smaller than house flies, fruit flies is composed of a variety of species, but how to handle it, and produce nearly the same destruction. Efforts to reduce and prevent fruit flies done by keeping the orchard, fruit falling in a vat containing input petisida, buried as deep as 75 cm and the soil surrounding the dismantled in order pupae hidden in the ground out, with oil-containing compound citranela isogenol metly eugenol and amyl alcohol which can bind the male fruit flies. One of the fly *Drosophila melanogaster* is *Drosophilidae* familia, these flies are small, at high populations can threaten human health, because of his interest in fruits and vegetables, especially materials that undergo fermentation (Matthews and Matthews, 1978). Imago of flies, measuring length, 2.5 mm - 4.0 mm (Charles and Noerman, 2005) is usually brownish yellow and brownish black. Melatakan eggs a female

can average 50-70 points per day eating an appropriate medium for the larvae is overly ripe fruit and vegetables that have been and will rot or material that is undergoing fermentation. In the dry season production of fruits and vegetables decreases the impact on the population of *D. melanogaster* fruit flies, fly *D. melanogaster* is one of the birds' feed (predator) populations of *D. melanogaster* in order to maintain the waste dry season rice bran (bran) The fermentasi for the development of breeding populations of *D. melanogaster*. The presence of predators in the world of agriculture is a free-living organism consuming, killing and devouring other organisms directly to the beneficial insects are not enough predators who can be identified from birds to microorganisms that can be separated on the farm. Based on the above meal authors chose to mengidentifikasi, populations of *D. melanogaster* and then developed through the artificial medium of rice bran (bran) fermented *Sacharomyces cerevoceae*, where applications and simple technology for the high economic value of feed bird that can improve the quality as well as the bird's nest swallow. This study aims to determine the populations of *D. melanogaster* medium banana shoes, which is compared with the artificial media. Followed by chemical analysis of *D. melanogaster*, protein, carbohydrates, fats, minerals and caloric value and determine the availability of feed to the birds' populations and determine the chemical compounds in the bird's nest covering nutrition, weight, thickness, proteins, carbohydrates, fats, minerals.

## B. METHODS

This research was conducted in the District Kamangta Tombulu Minahasa for 12 months. In locations with a height of 550 meters above sea level (asl) with an average temperature range of 26°C-30°C. Materials and Equipment Materials used in this study were rice bran / bran, bananas, shoes, yeast (*Sacharomyces cereviceae*), and honey. While the tool used was the fermentor, pH meter, hand refraktometer, petridish, microscope, auto Clav, mikrokeldhal micro, soklet, bomb calorimeter, AAS, and tools that support the analysis beaker *D. melanogaster*. The design of experiments The study was designed as a complete randomized trials with repeated three times for media banana shoes, as well as five treatments and three times re-made for the media. The parameters measured by pH, percent sucrose, determining population size medium banana shoes, the media and media metamorphosis in artificial insects are eaten by the percentage of birds', the chemical analysis of *D. melanogaster*, (proteins, fats, carbohydrates, minerals, and caloric value) body weight and feed the birds, the thickness and weight of the nest and the nest of proteins, carbohydrates, fats, minerals from bird nest feed treated *D. melanogaster*. Working procedure Before developing *D. melanogaster*, do pembedakan on swiftlet analyze the number of insects in the gizzard. Further develop *D. melanogaster* in cultured banana shoes, determine the pH and percent sucrose as a medium life cycle of egg, larva, pupa and imago. Moving the *D. melanogaster* of media banana shoes, artificial media to A, B, C, D, E, after the adult insects analyzed proteins, carbohydrates, fats, minerals, and the caloric value and determine the best medium of artificial media A, B, C, D, E to *D. melanogaster* insect breeding. Doing hatching birds' and weighing the birds' weight and feed children who were given the adult insect *D. melanogaster* from the first day until able to fly for 45 days. Measuring the thickness of the nest, nest weight and analyzes of bird nest (proteins, fats, carbohydrates, minerals) as well as the time required for nest building of birds treated and non-treated. observation. Made observations on the life cycle of each treatment by calculating the population of *D. melanogaster* cultured banana shoes, (generation 1 s / d 3rd generation), long metamorphosis and do the same thing on artificial media. The next rare daily feeding, weighing and amount of feed consumed, accounted for 45 days. Nesting observation for 65 days in the last swallows building harvested weighed, measured the thickness and weight of chemically analyzed nest swiftlet nests treated and non-treated.



## C. RESULT AND DISCUSSION

### A. Chemical Analysis Media Alamia Before dimasukkan in fermentator

#### 1 ). The level of acidity (pH) Natural Foods


Based on the observation level of acidity ( pH ) of natural food banana shoes , before inclusion in fermentator is 6 . Development of *D. melanogaster* population started after fertilasi consisting of two periods of the eggs in the embryonic period when fertilasi until such time as the young larvae hatch from eggs more or less within 24 hours , at this time the larvae eat incessantly.

The second period is after the eggs hatch called postembriodik development consisting of stage larva, pupa , and imago .

Table 1 . Development of *D. melanogaster* populations of banana shoes on artificial food .

### D. Population Growth and Long Life of *D. melanogaster* on Artificial Food .

#### 1 ) . banana Shoes



#### Eggs

Based on the analysis of variance , artificial food very significant effect on the E ( 416 grains ) , A ( 396 points ) , C ( 402 grains ) , D ( 382 grains ) , of the ( 292 grains ) , high egg populations . The invention relates to an insect host by substrate color and odor of volatile oil ( Chapman , 1971 ) . The eggs produced by female insects to a couple of times ( during the egg stage ) have not much effect on the population because the egg is not moving , not eating , and do not proliferate . The eggs of *D. melanogaster* . Elliptical -shaped objects are small and are usually placed on the surface of the food . Adult females begin to lay eggs on the second day after becoming adult flies and increased in the week to females laying 50-75 eggs per day . Within 10 days maximum of approximately 400-500 pieces of eggs .

eggs ; *D. melanogaster* vetillin coated by a thin membrane that surrounds the cytoplasm on the outside and thin silvia dianteriornya two tanks . According to ( Borror , 1992 ) , was a very strong thin membrane called the chorion and coated the outside of the egg chorion . flyblow Based on the analysis of variance , the food was highly significant , medium A ( 317 head ) , B ( 233.6 tail ) , C ( 322 head ) , D ( 306 animals ) , E ( 369.3 tail ) . Types of food , water content ,

and cross-sectional area . Food grain size effect on the development of a type of searangga .

Larvae ; *D. melanogaster* white , shaped like a segmented worm with a mouth and brownish black , has a pair of spiracles are located on the anterior and posterior ends . *D. melanogaster* experienced a skin substitute ( instars ) and lasts for three times where the first instar ( I ) larvae after hatching , the second instar ( II ) the appearance of the teeth in the mouth to eat black preparation for the third instar ( III ) the preparation to form pupae and moving kepinggir - edge fermentator the place dry and then stopped moving Pupa. Based on the analysis of variance , artificial food very significant effect on pupal populations . Media A ( 237 head ) , B ( 233.6 tail ) , C ( 241.6 tail ) , D ( 229.8 tail ) , E ( 277.3 tail ) .

Pupa ; *D. melanogaster* occurs when larvae instar ( III ) brown body that contract ( shorten ) the cuticle be hard and pigmented called fourth instar ( IV ) is characterized by the Establishment wing head and foot pads . In the pupal stage larvae in a passive state ( inactive ) . In this time the larvae towards / into adult flies Imago. Based on the analysis of variance , significant artificial food . Reproduction insect nutritional decline due to low population and development is also influenced by the type and amount of food . Media A ( 166 head ) , B ( 175.33 tail ) , C ( 164.33 tail ) , D ( 166.33 tail ) , E ( 195.33 tail ) .

Imago or adult flies aged about 10 days after discharge from the pupa , the wing can not be grown and still a bit pale and after approximately 10 days 9 hours fly females to mate . E. The protein in insects *D. melanogaster*. In insects as well as other animals one of the major biochemical activity at the cellular level is the synthesis of proteins , where the characteristic features of insects holo metabola is a change in the form of larvae into the imago ( adult ) at the time of metamorphosis . Fat bodies of insects is a major organ of metabolic processes a wide variety of biochemical substances and their role is crucial , especially in stadia pradewasa . Insects holometabola , during the development of insect larvae fat body responsible for the synthesis of various proteins in insect hemolimfanya synthesized in the fat body of an average protein synthesis at high body fat at the start of the growth of larvae and then decreases . Protein hemolimfa stored as vesibel to transport lipids ( fats ) , carbohydrates , and hormones . Synthesis and release of specific protein hemolimfa by fat body from larvae that were growing and the disposal and storage of these proteins in the same tissue for further development is controlled by hormones .

For that we should not be surprised when the insect protein in insects *D. melanogaster* showed high numbers , the protein results in laboratory analysis around 55.19 % . In addition to high-protein insects are relatively low fat content and a clean source of food by eating plants / fruits fresh . Chemical analyst Table *D. melanogaster* No. Parameter Analysis Method of Analysis Results

- 1 Protein 55.19 % Makrojeldahl
- 2 6.75 % Fat Extraction
- 3 1.80 % Carbohydrates Hydrolysis
- 4 Sodium ( Na ) 3.55 ppm AAS
- 5 Potassium ( K ) 50.34 ppm AAS
- 6 Calories 288.71 cal / g calculation

Insect *D. melanogaster* high protein ( 55.19 % ) as the birds' feed

F. Presentation Given Insects In Bird Treatment and Non- Treatment .

Table . Percentage Insects On The Swallow gizzard Kamangta What Not Provided treatment .


Ordo	Famili	Presentase (%)
Diptera	Drosophilidae	45
	Phoridae	
Cleoptera	Tenebrionidae	20
Homoptera	Aleyrodidae	10
Isoptera	Kaloptermitidae	10
Lain-lain	-	15*

Table. Percentage Insects On The Swallow gizzard Kamangta treated.

Ordo	Famili	Presentase (%)
Diptera	Phoridae	75
	Psychodidae	
	Drosophilidae	
Coleoptera	Tenebrionidae	20
	Silvamide	
Lain-lain	-	5*

Note \*: other; insect was identified as damaged. Swiftlet treated, take the order diptera and coleoptera. Non-free treatment to fly and prey on the order of the Homoptera, Isoptera, diptera, coleoptera and other insects are not detected proving that the location of the bird cage and the ecosystem environment favorable to the development of swiftlet (predators). Observation of the relationship between the number of birds feeding and body weight in birds treated with variable amounts of feed (X) and weight ( $\hat{Y}$ )  $r = 0.9$  with the regression  $y = 1.43x + 1.67$  where price  $x$  0.02 ,  $t_{hit} = 26, 16 > t_{0.99} (43) = 1.17$  this means that there is a highly significant positive correlation between the amount of feed and weight.

Weight Charts Swallow Treatment and Without Treatment



Based on observations of the thickness of a bird's nest , nest weight with treatment and without treatment with the null hypothesis as follows :

Reject , if  $t_{hit} > t_{tab}$

To test this hypothesis , compared to the average thickness of a given bird nest treatment ( ) with a bird's nest without treatment ( ) during the experiment .

= 0.98 with  $S_1 = 0.0968$

= 0.62 with  $S_2 = 0.0887$


$S_2 = 0.00862$

$t_{hit} = 12.26$

For  $\alpha = 0:01$  ,  $t_{hit} = 12:26 > t_{0,99} (38) = 2.42$  . This means that the average thickness of the bird's nest in the treatment of very , markedly higher than the thickness of the bird's nest naturally ( without treatment ) .

The thickness and weight of the bird's nest is very real treatment is higher than the thickness of the non- treatment birds' nests  $\alpha = 0.01$  ,  $t_{hit} = 12.26 > t_{0.99} (38) = 2.42$  . Reject  $H_0$  if  $t_{hit} > t_{0.99} (38) = 2.42$ , this means that the average weight of edible bird's nest in a very real treatment is higher than the weight of non- treatment birds' nests .

Graph thickness Swallow's Nest Treatment and Without Treatment



#### Heavy Bird's Nest

Based on observations, it turns heavy bird nest treated higher than the natural bird's nest (without treatment), null hypothesis is formulated as:

Reject, if  $t_{hit} > t_{tab}$

To test this hypothesis, compared to the average weight of a given bird's nest treatment ( ) with a natural bird nest / no treatment ( ) during the experiment.

= 8.56 with  $S_1 = 0.4867$


= 7.52 with  $S_2 = 0.4588$

$S_2 = 0.2237$

(Vincent, 1991)

$t_{hit} = 6.95$  For  $\alpha = 0:01$ ,  $t_{hit} = 6.95 > t_{0.99} (38) = 2.42$ . This means that the average weight of birds' nests in the treatment of very , markedly higher than the natural bird's nest weight (without treatment).

Weight Charts Swallow's Nest Treatment and Without Treatment



When food available for the birds in the vicinity of the building energy treated birds to search for food, does not require a long time to find / hunt for food outside the building, especially in the dry season. When hunting birds feed away from buildings, insects gained less than the needs and birds suffering from fatigue, the possibility of getting lost or birds can not return original place of occupancy. Backup saliva prepared for nest building and in particular the protein is reduced impact on the formation of a nest of thin / thick not.

Table. Results of Chemical Analysis of swallow's nests

No	Parameter (%)	Analysis Results		Analysis Method
		Treatment	Without Treatment	
1	Fat	0,30	0,21	Hydrolysis
2	Protein	52,05	52,47	Makrokjeldhal
3	Carbohydrate	16,32	16,33	Hydrolysis
4	Calcium	2,26	0,19	AAS
5	Potassium	2,36	3,18	AAS
6	Magnesium	0,05	0,07	AAS
7	Sodium	0,13	0,18	AAS

CONCLUSION

Insect *D. melanogaster* is an insect living on fruits and media that undergo fermentation , it can be seen clearly when fruits or vegetables and garbage who started / would rot . These insects are interesting to study because living around humans ( insects settlement ) of the order Diptera Drosophilidae family .

These insects examined in the Village Kamangta with a height of 500-600 m above sea level in fermentator intended bred again swiftlet larvae feed children , imago ( adult ) for the adult birds' feed .

Some of the scientific information about the contribution of *D. melanogaster* is used to control feed swiftlets ( *collacalia fuchifaga* ) of larvae and adults very well with the value of 55.19 % protein , which is bred from natural media ( banana shoes ) and fermentation of bran / rice bran .

1 . Based on the analysis of variance showed that the natural medium of banana shoes gave a positive response in the growth of insect populations of *D. melanogaster* , for artificial feeding fermented bran treatment E to *D. melanogaster* best of feed nutrients banana shoe types significantly influenced the development of the fly population *D. melanogaster* fruit and the largest population in the natural medium banana shoes . Besides comparison nisba kelamin1 male and 5 females and *D. melanogaster* population growth process starts from mating behavior , egg laying , and the process of metamorphosis . 2 . Artificial feed ( artificial food ) intended to divert the insects *D. melanogaster* from multiplying in the months naturally cultivated , while the bran / rice bran fermented pest activity is expected to result from *D. melanogaster* decreases and reduced and will be a source of feed swiftlets ( predators ) . Predators can be cultivated in areas mountains at an altitude up to 1000m above sea level between 300m asl apart in coastal areas .

3 . Feed type ( insect *D. melanogaster* ) as the birds' feed high protein significantly affect the thickness of the bird's nest . For  $\alpha = 0:01$  ,  $t_{hit} = 6.95 > t_{0.99} ( 38 ) = 2.42$  and weight of the non- treatment nest .

In organoleptic form nests did not differ either in form and color , and the provision of *D. melanogaster* as feed can reduce the duration of post -harvest dry season from 80 days to 65 days .

4 . There is a linear relationship between the amount of feed and weight of children swallow , for  $\alpha = 26.16$   $00:01$   $t_{hit} > t_0$  ,  $95 ( 43 ) = 1.67$  means that there is a highly significant positive correlation between the amount of feed ( x ) and weight ( y ) the regression  $Y = 1.43 x + 1.67$  . Imago *D. melanogaster* , have a high protein 55.19 % for  $\alpha = 4.56$   $0:01$   $t_{hit} > t_0$  ,  $95 ( 43 ) = 1.67$  weight chicks fed researcher significantly higher than the child's weight bird that fed their parent in other words that the *D. melanogaster* both being fed to birds' . B. Advice This study can be used as a reference for the control of insect pests of fruit fly *D. melanogaster* by exploiting predators ( birds' ) is very beneficial because it targets precisely to the target pest , and does not provide impact resistance on non-target insects than the use of pesticides . Another advantage is the predator birds' nests can produce high economic value of birds and insects *D. melanogaster* easy to breed in a short time (10-13 days ) , in addition to the insect pest is below 4 mm , which can lead to pests and diseases in plants and plants can be controlled without the use of pesticides . Voting should be based on the principle of sustainability in order to nest swiftlet population does not experience stress and nest remains sustainable production in post-harvest production , as well as the quality and quantity is maintained nest .

1	1.0	2.5	1.0
2	1.5	3.5	1.5
3	2.0	4.5	2.0
4	2.5	5.5	2.5
5	3.0	6.5	3.0