

The bioactive contents and
antioxidant activity of honey
bee nest extract of *Apis dorsata*
Binghami from the North
Sulawesi
by Mocosuli Yeremia

Submission date: 27-Mar-2023 10:40AM (UTC+0700)

Submission ID: 2047575285

File name: document_17.pdf (401.66K)

Word count: 5838

Character count: 31584

The bioactive contents and antioxidant activity of honey bee nest extract of *Apis dorsata* Binghami from the North Sulawesi

Mokosuli Yermia Samuel^{1,2*}, Eva Sherly Nonke Kaunang³, Jacklin Stella Manopo²

¹Laboratory of Bioactivity and Molecular Biology, State University of Manado, Kampus Unima Tondano, Minahasa, Sulawesi Utara, Indonesia

²Departement of Biology, Faculty Mathematics and Natural Science, State University of Manado, Kampus Unima Tondano, Minahasa, Sulawesi Utara, Indonesia.

³Departement of Education Biology, Faculty Mathematics and Natural Science, State University of Manado, Kampus Unima Tondano, Minahasa, Sulawesi Utara, Indonesia

*Corresponding author email: yermiamokosuli@unima.ac.id

Received March 8; 2019; Accepted May 20; 2019; Available online November 30, 2019

ABSTRACT. *Apis dorsata* Binghami is a Sulawesi endemic honey bee. *Apis dorsata* Binghami cannot be bred, so it still lives wildly in the forests of Sulawesi. However, *Apis dorsata* Binghami produces more honey, compared to all honey bee species. Furthermore, the diversity of plants as a source of nectar, pollen and plant resin, which is used in the formation of nests and honey is more, than all types of honey bees in the world. Ethnomedically, the Minahasa community has long used honeynest for degenerative diseases such as hyperlipidemia and cancer. Nevertheless, there have been no research reports on bioactive content and bioactivity of *Apis dorsata* Binghami nest extract. This research was aimed to determine the bioactive content of honey bee nest and to obtain the inhibitory concentration 50 (IC₅₀) antioxidant activity of honey bee nest extract of *Apis dorsata* Binghami. Honey bee nest was obtained directly from the forest of Minahasa peninsula, North Sulawesi, Indonesia. Extraction of fresh honey bee nest was conducted using maceration method. Bioactive content analysis was carried out by the Harborne method, followed by analysis using UV Vis spectrophotometer and High performance liquid chromatography. IC₅₀ antioxidant activity of honey bee nest extract was obtained using DPPH free radical reduction method. The results showed that *Apis dorsata* Binghami honey bee nest extract contained alkaloids, flavonoids, saponins, tannins, steroids and triterpenoids. Identified flavonoids displayed the highest phytochemical content. Based on the results of HPLC and UV Vis spectrophotometer analysis, there were 20 flavonoid derivatives found in honey bee nest samples in Minahasa. Ethanol extract and n-hexane extract showed high free radical reduction activity compared to vitamin C as a control treatment. However, ethanol extract produced the highest DPPH free radical reduction activity.

Keywords : Bioactive, *Apis dorsata* Binghami, nest extract, antioxidants

INTRODUCTION

Apis dorsata Binghami is an endemic bee of Sulawesi that still lives naturally in the forest and cannot yet be bred. *Apis dorsata* Binghami was discovered by Alfred Russel Wallace during an expedition to Sulawesi Island (Celebes) in the 18th century (Hadisoesilo, 2001; Raffiudin, 2002; Mokosuli, 2013). *Apis dorsata* Binghami produces the most productive honey compared to other honeybee species (Otis, 1991; Hadisoesilo, 2001). This bee makes a nest with only one comb hanging from tree trunks and twigs, open ceilings, and rocky cliffs (Hadisoesilo, 2001; Raffiudin, 2002). In spite of these advantages, they are not commonly cultivated as *Apis mellifera* and *Apis cerana* honeybees. Until now, scientists have not succeeded in cultivating *A. dorsata* in a closed form, so that it still lives naturally in the forest (Otis 1991; Hadisoesilo 2001; Raffiudin, 2002). However, deforestation is rapidly decreasing

their nesting in Minahasa, North Sulawesi, Indonesia (Mokosuli, 2013; Hadisoesilo et al. 2007; Sawaya et al. 2009), affecting their migration and survival. Therefore, information on the composition of *Apis dorsata* Binghami nest and propolis of this native honeybees, as well as the plants they visit as source of nectar, pollen, resins and other compounds from plants, is of primary importance.

Honey, bee venom and honey bee nest are honey bee products, which are prepared by worker bees. The composition and content of honey, venom and honey bee nest is very dependent on the diversity of plant sources of bee feed. Unlike *Apis mellifera*, where the diversity of food sources is limited, *Apis dorsata* Binghami has a very large variety of feed sources. The range of flying worker bees in search of feed plants is also further than *Apis mellifera* because of differences in body size (Mokosuli, 2013). *Apis dorsata* is known as a giant bee because it has the

largest body size compared to other honey bee species on earth (Hadisoesoilo 2001; Raffiudin, 2002). Previous research has been conducted, found that the venom content and *Apis dorsata* Binghami honey are strongly influenced by variations of flowering plants adjacent to their nests. *Apis dorsata* found in the Kombi of Minahasa Regency in the season of clove plants produces honey with the aroma of cloves (Mokosuli, 2013). In addition to honey and venom, honey bee nest is a potential bioactive source of medicine that has the potential to be studied.

The honey bee nest is the structure used by bees as a place to live and raise their offspring. The inside of the honey bee nest is a collection of hexagonal structures made of wax. Bees use this hexagonal space to store honey, bee pollen, eggs, larvae and bee pupae (Nazzi, 2016). The honey bee nest is a refuge for bee colonies from attacks by bacteria, fungi, viruses and predators, as well as a place to produce honey, bee pollen, and a place to grow bee eggs. The condition of the honey bee nest greatly affects the quality of honey and what is produced. Honey is generally free from pathogenic microbes (Perez, Suarez, Pena-Vera, onzalez & Vit, 2013). The content of compounds in honey bee nest serves as a protective and determinant of the quality of honey including flavonoids which are natural phenol compounds and bees wax (Ra'ed, Ibrahim, Rula & Mosa, 2008; Pasupuleti, Sammugam, Ramesh & Gan, 2017).

Propolis is the largest component of honey bee nest constituent. Propolis is a sticky, gummy, resinous substance collected by honeybees from various plant sources (Coneac et al, 2008). Propolis is produced by bees from sap taken from parts of plants that produce sap, especially plant buds (Nazzi, 2016; Bankova, 2005). The sap is the basic ingredient for propolis. Sap is brought into the honey bee nest by worker bees and mixed with wax (a type of wax) and flower pollen. With the help of bee saliva, this mixture is processed to become flexible to form propolis (Suseno, 2009). Propolis contains aromatic compounds, flavonoids, quercetin, terpenoids, and sugar. In addition, there are also minerals Fe, Ca, Mg, K, Na, and Zn. Natural propolis contains a number of amino acids such as valine, isoleucine, leucine, proline, alanine, and glycine which play a role in the formation of body cells. Propolis is also rich in vitamin B1, vitamin B2, vitamin B3, and vitamin B6 (Choi, Shimomura, Kumazawa & Ahn, 2013).

Bee propolis extract has antidiabetic activity, antiaterogenic, antimicrobial and antifungal mainly due to polyphenol content (Daleprane & Abdalla, 2013; Barrientos et al. 2013; Saavedra et al. 2011; Veloz et al. 2016). The benefits of flavonoids in the

human body function as antioxidants so it is very good for cancer prevention. The benefits of flavonoids include anti-inflammatory, preventing bone loss, and as an antibiotic (anti-bacterial and anti-virus) (Waji and Sugrani, 2009; Redha, 2010; Seal, 2016). The benefits of quercetin are believed to protect the body from several types of degenerative diseases by preventing the occurrence of fat peroxidation (Waji & Sugrani, 2009). Quercetin shows the ability to prevent the oxidation process of LDL cholesterol by capturing free radicals and holding transition metal ions (Repi, R. A., Mokosuli, Y.S., Ngangi, J., & Sumipouw, H. M. 2013). Polyphenol derivatives as antioxidants can stabilize free radicals by complementing the lack of electrons possessed by free radicals, and inhibiting the occurrence of chain reactions of free radical formation (Mokosuli, 2008). Polyphenols are components responsible for antioxidant activity in fruits and vegetables (Hattenschwiller and Vitousek 2000). Aside from being an antioxidant, polyphenols can also reduce cholesterol, LDL and triglyceride levels. The reduction mechanism is by increasing the activity of lipoprotein lipase, so that the catabolism of triglyceride-rich lipoproteins such as VLDL and IDL increases (Redha, 2010). HDL cholesterol levels increase indirectly due to decreased triglyceride levels of VLDL or due to increased production of apo AI and apo AII (Inggrid & Santoso, 2014). The effect of decreasing LDL cholesterol is thought to be related to increased clearance of VLDL and IDL in the liver so that LDL production decreases (Mokosuli, 2013). Ethnomedically, the Minahasa community has long used honeynest for degenerative diseases such as hyperlipidemia and cancer. Nevertheless, there have been no research reports on bioactive content and bioactivity of *Apis dorsata* Binghami nest extract.

EXPERIMENTAL SECTION

Tools and Materials

The tools used in this study include : UV-Vis Spectrophotometer *Parkin Elmer*, Nanophotometer *Implan*, Rotary Evaporator *Heidolph*, Automatic Qiaexel electrophoresis *Qiagen*, HPLC Shimizu, micropipette eppendorf, sample boxes, waterbaths, UV Cabinets, glassware etc. The materials used include: ethanol p.a. merck, Methanol p.a. Brand, Merck n-hexane, Merck ethyl acetate, Merck chloroform, Pareaksi Dragendorf, Pareaksi Mayer, Pareaksi Wagner, HCL, Diphenyl pikri hydrazill (DPPH) Merck, propylene glycol (Merck) $K_2Cr_2O_7$, 1 N, H_2SO_4 , acetonitrile, standard phenol (Merck), standard flavonoids (Merck), antihyperlipidemia analysis kits, bioactive analysis kits, HPLC columns etc.




Figure 1. Location of a honey bee nest samples (red white circle). (Map source: google earth, www.google.com)

Honey Bee Nest Samples

Honey bee nest were obtained from the Minahasa peninsula, North Sulawesi province, Indonesia (**Figure 1**). Honey bee nest is taken at natural nesting sites in the forest, collected in a sample box with controlled temperature (temperature 25 °C). The sample was carried out to the Laboratory Bioactivity and Biomolecular Department of Biology, State University of Manado, for analysis.

Extraction

Extraction of *Apis dorsata* Binghami honey bee nest, carried out by maceration method. The fresh honey bee nest consists of a bag of honey, a pollen bag, an egg bag, and a nest cover. Whole honey bee nest, then blended until smooth. Comparison of honey bee nest solvents and simplicia 1: 4 (b / v), (250 grams of honey bee nest powder macerated with 1000 ml of 70% ethanol), at room temperature. After 48 hours, screening with Whatman number 4 filter paper was carried out. Solvents on the filtrate were identified with a Heidolp rotary evaporator, at a temperature of 40°C, 55 rpm. The extract obtained is then called honey bee nest ethanol extract. While the residue continued with maseration using n-hexane as much as 1000 ml. After 48 hours, screening was done with Whatman no. 4 filter paper. Solvents were evaporated with a Heidolp rotary evaporator at 40°C, 55 rpm. The extract obtained referred to as honey bee nest n-hexane extract.

Bioactive Content Analysis

Phytochemical analysis

Content analysis of phytochemical groups using the method of Harborne (2008) and UV-Vis Spectrophotometer. The crude honey bee nest extract was tested using Wagner solution, Dragendorff solution and Meyer solution. The components analyzed include the content of Alkaloids, Flavonoids, Saponins, Tannins, Streorids and Triterpenoids.

Bioactive analysis of the HPLC method

Flavonoid analysis of Honey Bee nest was conducted using HPLC. A sample of 2 g honey bee nest plus 14 mL acetonitril 70%, then allowed to stand for 24 hours. Samples were filtered using filter paper (Whatmann No.41) and PVDF filter (Milipore). Measurement of phenol levels were performed using HPLC.

RESULTS AND DISCUSSION

Extraction

Extraction is the initial stage to isolate potentially drug-active compounds from plant or animal simplicia. The most important factor, influencing the extraction results, namely solvent, time, and temperature in extraction (Yang, Wang, Ke, Jiang & Ying, 2007). *Apis dorsata* Binghami honey bee nest, obtained from the Minahasa forest, was freshly prepared for extraction (**Figure 1**). *Apis dorsata* Binghami honey bee nest extraction was carried out using 70% ethanol solvent, and n-hexane. The honey bee nest is freshly blended, into powder, then as much as 250 g, the powder is macerated with 800 mL 70% ethanol, for 48 hours at room temperature. The solvent contained in the filtrate was evaporated used a rotary evaporator, at a temperature of 40 °C, at 48-50 rpm, then called ethanol extract. The residue was again macerated with 800 mL n-hexane for 2 x 24 hours at room temperature. The maceration results are then filtered to obtain the filtrate. The solvent was evaporated with a rotary evaporator at a temperature of 40 °C, 50 rpm. The extraction results are then called n-hexane extracts (**Table 1**). Ethanol extract is brown with a distinctive honey bee nest odor, while n-hexane extract is yellow with a distinctive honey bee nest odor (**Figure 1**).

Table 1. Yields of *Apis dorsata* Binghami honey bee nest extract

Samples	Solvents	Samples Weight (g)	Yields (%)	Extract Weight (g)	Colours
A1	Ethanol 70%	200	7.75	5.32	Brown
	n-heksan	200	8.45	6.45	Yellow
A2	Ethanol 70%	200	8.53	5.45	Brown
	n-heksan	200	8.91	6.53	Yellow



Figure 2. (a). *Apis dorsata* Binghami honey bee nest and (b). *Apis dorsata* Binghami honey bee nest extract

Tabel 2. Phytochemicals analysis results

Group of Phytochemistry Compounds	Results			
	A1Et.OH	A1nh	A2EtOH	A2nh
Alkaloid	+	++	++	+
Flavonoid	++	+	++	+
Saponin	++	+	++	++
Tanin	+	-	+	+
Triterpenoid	+	+	+	+
Steroid	++	++	+	++

Description :

(+) sign indicates the level of color intensity and deposits formed, after being given the test reaction. EtOH (ethanol extract), nh (n-hexane extract).

The most widely used method in the study of plant materials as a bioactive source for medicine is the maceration method. In addition to the maceration method, percolation methods, soxhlet and steam distillation are also often used. Nevertheless, percolation methods are only good for soluble organic compounds, while steam and steam distillation are only good for heat-resistant compounds (Faraouq, 2003; Lenny, 2006; Mokosuli, 2008). According to Faraouq (2003), the extraction of simplicia for the purpose of herbal medicine is the best use of ethanol solvents. Ethanol can mix with water in various comparisons, and it is easy to evaporate the residue in the extract. Methanol, ethylacetate or hexane solvents are not allowed because of toxic residues produced (BPOM, 2018). According to Trusheva, Bankova & Trunkova, (2007), extracting propolis with ethanol by comparing several extraction methods namely maceration, EU (ultrasound extraction) and MAE (microwave assisted extraction), it turned out that the maceration method produced a total yield of 55.58% greater than the EU and MAE methods with each yield was 41% and 53%. The highest yield percentage was obtained

from ethanol extract both on A1 and A2, therefore the two extracts were continued in an antioxidant activity bioassay.

Phytochemical Analysis

Screening content of phytochemicals is a standard procedure in the exploration of plant and / or animal extracts, for the purpose of testing drug activity. Phytochemical analysis aims to determine secondary metabolites, which are contained in honey bee nest extract. Honey bee nest is formed, among others, from the sap of plants taken by worker bees. In A1 ethanol extract, it was identified that all classes of secondary metabolites were tested. Flavonoids, Saponins and Steroids are found in higher intensity than other secondary metabolites. While n-hexane extract obtained alkaloid and steroid content in high intensity. In the ethanol extract A2, obtained the content of alkaloids, flavonoids and saponins with a higher intensity than other phytochemicals tested. n-hexane A2 extract showed saponin content and steroids with higher intensity than other phytochemical groups (Table 2).




Figure 3. Spectrophotograph of ethanol extract of *Apis dorsata* Binghami honey bee nest




Figure 4. Spectrophotograph of n-hexan extract of *Apis dorsata* Binghami honey bee nest

In forming a nest, *Apis dorsata* Binghami requires more than 100 plants, a source of sap (Hadisoesoilo 2001; Raffiudin, 2002). The latex is used in forming hexagonal spaces in the honey bee nest, where bee eggs are placed. Previous research, *Apis dorsata* Binghami nests in Minahasa have specific characteristics (Mokosuli, 2013). Some of the plants that are visited by bees are *Syzygium aromaticum*, *Ficus* sp., *Ficus minahassae*, *Durio zebethinus*, *Lansium minahassae*, *Elemerelia celebica* L, *Mangifera indica*, *Cocos nucifera*, *Garcinia magostana*, *Nephelium lappaceum* etc.

This reinforces the results of research that *Apis dorsata* Binghami bee nest contains all phytochemical classes. Extraction is done by maceration and fractionation methods. Ethanol solvents have two different polarity groups, namely polar hydroxyl groups and nonpolar alkyl groups. The existence of this cluster so that compounds with different polarity levels will be extracted in ethanol. n-hexane solvents are non-polar, this causes the secondary non-polar metabolites to be properly


withdrawn. Profile of the UV Vis spectrophotometer, both extracts are shown in **Figure 3** and **Figure 4**.

Flavonoid analysis with HPLC

Analysis of the content of flavonoid compounds was carried out using High performance liquid chromatography. HPLC can separate compounds from the mixture in the honey bee nest extract. Separation of compounds is known in retention time (Seal, 2016). Retention time is the time needed for the compound to move, when it is first injected into the chromatography, to the maximum peak measured (Seal, 2016). Unknown compounds can be observed well using UV light at wavelengths between 200-220 nm (close to the final absorbance), this is due to the fact that organic compounds have absorbance at the spectrum distance (De Oliveira, Nakashima, De Souza & Frehse, 2001; Rosli et al., 2016). Based on retention, there are 35 types of active compounds in A1 honey bee nest extract. Three compounds were found with a percentage above 10%. First, the active compound content was

23.63% with a retention time of 2.773 minutes. Second, the content of the active compound was 14.28% with a retention time of 3.793 minutes, and the third content of the compound was 11.88% with a retention time of 2.003 minutes. Quercetin standard shows retention time of 11.854 (Figure 3). Quercetin

was identified in A1 extract with a higher content compared to extract A2. The lower the sample concentration, the higher the retention time value. This causes the compound to be more difficult to separate from other compounds, if the levels in the sample are not enough.




Retention time A1 (EtOH) extract

UV-VIS Results

Retention Time	Area	Area %	Height	Height %
1.580	111790	2.24	12152	2.29
2.033	319951	6.42	63113	11.88
2.160	137408	2.76	22272	4.19
2.350	132494	2.66	13665	2.57
2.530	91519	1.48	13768	2.59
2.650	153338	3.08	28984	5.46
2.773	920666	18.47	125550	23.63
3.010	80798	1.62	15293	2.88
3.153	375551	7.53	36147	6.80
3.587	121473	2.44	15063	2.84
3.793	673915	13.52	75838	14.28
4.363	159673	3.20	15135	2.85
4.687	136630	2.74	15020	2.83
5.337	320943	6.44	20957	3.94
5.907	57300	1.15	4851	0.91
6.317	94395	1.89	4208	0.79
6.690	54456	1.09	3691	0.69
7.800	150424	3.02	7021	1.32
8.593	60223	1.21	3396	0.64
8.893	115809	2.32	5344	1.01
10.810	26331	0.53	1346	0.25
11.267	19501	0.39	1061	0.20
11.883	29545	0.59	1266	0.24
13.400	244190	4.90	7183	1.35
14.657	126413	2.54	3214	0.60
15.017	10649	0.21	2699	0.51
15.157	30248	0.61	2777	0.52
15.300	61692	1.24	2775	0.52
16.473	37909	0.76	1135	0.21
17.983	27225	0.55	1165	0.22
18.127	30895	0.62	1218	0.23
20.733	37102	0.74	1547	0.29
20.850	23213	0.47	1576	0.30
21.203	11130	0.22	820	0.15


Figure 5. Results of UV Vis Spectrophotometry analysis of A1 extract after analysis by HPLC



UV-VIS Results

Retention Time	Area	Area %	Heigth	Heigth %
1.443	20835	14.46	2570	13.99
1.557	49079	34.06	5207	28.35
2.030	64071	44.47	9974	54.30
2.710	10104	7.01	619	3.37
Total	144089	100.00	18370	100.00

Retention time A2 (*n*-hexane) extract



UV-VIS Results

Retention Time	Area	Area %	Heigth	Heigth %
1.607	67545	3.67	3419	5.25
1.997	27648	1.50	2902	4.45
11.853	1743535	94.82	58835	90.30

Figure 6. Quercetin standard retention time and results of UV Vis Spectrophotometer analysis




Figure 7. Extract inhibition activity at various test concentrations of DPPH free radicals

Test of antioxidant activity

Antioxidant activity test method DPPH 1,1-diphenyl-2-picrylhydrazyl is widely used to determine the antioxidant potential of plant extracts. Antioxidant activity in honey bee nest extract, reacts with DPPH free radicals by donating hydrogen atoms, causes changes in DPPH color from purple to yellow. Color intensity was measured by spectrophotometer at a wavelength of 517 nm (Mokosuli 2008). The principle of this antioxidant activity test method is to quantitatively measure antioxidant activity by measuring DPPH radical capture by a compound that has antioxidant activity using UV-Vis spectrophotometry, so that the value of activity of free radical reduction is expressed as IC₅₀ value (Inhibitory Concentration) (Kikuzaki & Nakatani, 1993). Antioxidant activity was obtained from the percentage of inhibition of *Apis dorsata* Binghami honey bee nest extract, against DPPH radical. The percentage of inhibition was obtained from differences in absorption between DPPH absorbance and absorbance of the sample measured by UV-Vis spectrophotometer. The amount of antioxidant activity is indicated by the IC₅₀ value, which is the concentration of the sample solution needed to inhibit 50% DPPH free radicals. Based on the distribution of test concentrations, ethanol extract (Aetoh) had the best free radical inhibitory percent compared to n-hexane (Anh) extract (Figure 7).

Table 4. IC₅₀ value of the antioxidant activity of *Apis dorsata* Binghami honey bee nest extract

Samples	IC ₅₀ (ppm)
Vit C (control)	6.73
Ethanol extract (AEtOH)	6.690
n-heksan extract (Anh)	6.766

Compared to the positive control of vitamin C with IC₅₀ (6.73 mg / L), ethanol extract has a better

IC₅₀ value of 6.69 mg / L. Whereas n-hexane extract has a lower IC₅₀ value of 6.76 mg/l (Table 1). Based on the IC₅₀ obtained, honey bee nest extract is very potential to be used as a source of bioactive antioxidants. The honey bee nest extract used is still a crude extract while vitamin C used is a control is a pure compound. A compound is said to be a very strong antioxidant if the IC₅₀ value is <50 ppm, strong for IC₅₀ is 50-100 ppm, while if IC₅₀ is 100-150 ppm, it is weak if IC₅₀ is 150-200 ppm, and very weak if the IC₅₀ value is > 200 ppm (Mardawati, Filianty & Marta, (2008)

DPPH free radical reduction activity of ethanol extract was stronger than n-hexane extract, supported by the content of active compounds contained in the extract. Based on the analysis of the content of phytochemical groups, ethanol extract has higher intensity of flavonoid, saponin and tannin content compared to n-hexane extract. The group of phenolic compounds, has been widely reported to have strong antioxidant activity. The n-hexane extract contains more compounds with lipid basic precursors such as triterpenoids and steroids.

Flavonoids act as free radical scavenger by giving hydrogen atoms to free radicals (Sanomiya, Fonseca & Silva, 2005). Free radical scavenging activity of flavonoids depends on the molecular structure and the form of substitution of hydroxyl groups such as the ability of phenolic hydrogen and the possibility of stabilization by phenoxyl radicals through hydrogen bonds or electron delocalisation. Structural activity activities (SAR) of important flavonoids are known, namely the number and location of OH phenolic groups that play a role in neutralizing free radicals (Amic, Davidovic, Beslo, & Trinajstic, 2003). Based on the analysis by HPLC there were more than 20 types of flavonoids contained in the ethanol extract of *Apis dorsata* Binghami honey bee nest while in n-hexane extract only found 4 flavonoid compounds. Flavonoid compounds such as quercetin, kaempferol,

myricetin, apigenin, luteolin, vitexin and isovitexin are found in cereals, vegetables, fruits and their processed products with various ingredients and most have antioxidant properties. This has reinforced the notion that flavonoids have certain biological effects, related to their antioxidative properties.

A mixture of several polyphenol compounds is able to function synergistically with other components as antioxidants and damping free radicals and prevention of various diseases (Meskin, Bidlack, Davies & Omaye, 2002). However, the antioxidant activity testing method is very influential in producing IC50 values (Shalaby and Sanaa, 2012). The difference in the value of anti-oxidant activity is caused by the extraction method, the testing method and the operating conditions used when the extraction process also differs (solvent volume, leaf powder size, extraction time, temperature, and pressure). Antioxidant activity is influenced by extraction methods and operating conditions used during extraction (Chuu et al. 2012).

CONCLUSIONS

Apis dorsata Binghami honey bee nest extract from Minahasa based on phytochemical analysis was identified to contain alkaloids, flavonoids, saponins, tannins, steroids and triterpenoids. Identified flavonoids are the highest phytochemical content. Based on the results of HPLC and UV Vis spectrophotometer analysis, there were 20 flavonoid derivatives found in honey bee nest samples in Minahasa. Ethanol extract and n-hexane extract showed high free radical reduction activity compared to vitamin C control. However, ethanol extract was found to have the highest DPPH free radical reduction activity.

Recommendations

In the future, it is necessary to study anticancer and antihyperlipidemia activities of *Apis dorsata* Binghami nest extract.

ACKNOWLEDGMENTS

Gratitude to the directorate of research and community service, Ministry of research, technology and higher education, Republic of Indonesia, which has financed the implementation of this research, in 2018.

REFERENCES

- Amic, D., Davidovic-Amic, D., Beslo, D., Trinajstić, N. (2003). Structure-Radical Scavenging Activity Relationships of Flavonoids. *Croatica Chemica Acta*. 76(1), 55-61.
- Bankova, V. 2005. Recent trends and important developments in propolis research. *eCAM*;2(1), 29–32. doi:10.1093/ecam/neh059
- Barrientos, L., Herrera, C. L., Montenegro, G., Ortega, X., Veloz, J., Alvear, M., Cuevas, A., Ovedra, N., & Salazar, L.A. (2013). Chemical and botanical characterization of Chilean propolis and biological activity on cariogenic bacteria *Streptococcus mutans* and *Streptococcus sobrinus*. *Brazilian Journal of Microbiology* 44(2), 577-585. <http://dx.doi.org/10.1590/S1517-83822013000200038>.
- B POM R. I. (Badan Pengawasan Obat dan Makanan Republik Indonesia). (2018). Peraturan badan pengawas obat dan makanan republik indonesia tentang kriteria dan tata laksana ekstraksi obat tradisional. <http://jdih.pom.go.id/rancanganpublikshowpdf.php?u>.
- Choi, S. J., Shimomura, K., Kumazawa, S., & Ahn, M. R. (2013). Antioxidant Properties and Phenolic Composition of Propolis from Various Geographic Regions in Korea. *Food Science and Technology Research*, 19 (2), 211–222. <https://doi.org/10.3136/fstr.19.211>
- Chuu, C. P., Lin H. P., Ciaccio, M. F., Kokontis J. M., Hause R. J., Hiipakka R. A., Liao, S., & Jones R. B. (2012). Caffeic Acid Phenethyl Ester Suppresses the Proliferation of Human Prostate Cancer Cells through Inhibition of p70S6K and Akt Signaling Networks. *Cancer Prevention Research*,; 5 (5): 788 DOI: 10.1158/1940-6207.CAPR-12-0004-T.
- Coneac, G., Gatifanu, E., Hadaruga, D. I., Hadaruga, N. G., Pinzaru, I. A., & Bandur, G. (2008). Flavonoid Content of Propolis from the West Side of Romania and Correlation with the Antioxidant Activity. *Chemical Bulletin Politechn* 53(67):56-60.
- De Oliveira, B. H., Nakashima, T., De Souza Filho, J. D., & Frehse, F. L. (2001). HPLC Analysis of Flavonoids in *Eupatorium littorale*. *Journal of the Brazilian Chemical Society*, 12(2), 243–246. <https://doi.org/10.1590/S0103-50532001000200019>
- Daleprane, J. B., & Abdall, D. S. (2013). Emerging Roles of Propolis: Antioxidant, Cardioprotective, and Antiangiogenic Actions. *Evidence-Based Complementary and Alternative Medicine*. Volume 2013, Article ID 175135, 8 pages (1-8) <http://dx.doi.org/10.1155/2013/175135>
- Farooqui, T., and Farooqui, A. A. (2010). Molecular Mechanism Underlying the Therapeutic Activities of Propolis: A Critical Review. *Current Nutrition & Food Science*, pp.1-15, Vol. 6, No.3
- Farouq, (2003). Ekstrak sebagai salah satu pengembangan bentuk obat tradisional. *Prosiding Seminar Nasional Tumbuhan Obat Indonesia XXIII, Jakarta*. 45-52

- 46 Hadisoesilo, S. (2001). Review : Keanekaragaman Spesies Lebah Madu Asli Indonesia. *Biodiversitas* 2, 123–128.
- 2 Hadisoesilo, S., Raffiudin, R., Susanti, W., Atmowidi, T., Hepburn C, Radloff S. E., Fuchs S., & Hepburn R. (2007). Morphometric analysis and biogeography of *Apis koschevnikovi* Enderlein (1906). *Apidologie* 39 DOI: 10.1051/apido:2008029
- 31 Harborne, J. B. (2008). Metode Fitokimia, penuntun dan cara modern menganalisis tumbuhanPenerjemah : Padmawinata K dan Soediro I. Penerbit Bandung
- Hattenschwiler, S., & Vitousek, P. M. (2000). The role of polyphenols in terrestrial ecosystem nutrient cycling. *Trends Ecology Evolution*. Jun;15(6):238-243.
- 24 Ingggrid, M., & Santoso, H. (2014). Ekstraksi Antioksidan Dan Senyawa Aktif Dari Buah Kiwi (*Actinidia deliciosa*). *Lembaga Penelitian Dan Pengabdian Kepada Masyarakat*, III(3), 43.
- Kikuzaki, H., & Nakatani, N. (1993). Antioxidant effects of some ginger constituents. *Journal of Food Science* 58:1407-10.
- 32 Lenny, S. (2006). Isolasi dan uji bioaktivitas kandungan kimia utama puding merah dengan metode uji brine shrimp. USU Repository, Medan.
- 17 Meskin, M. S., Bidlack W. R., Davies, A. J., & Omaye S. T. (2002). *Phytochemicals in Nutrition and Health*. CRC Press, London, New York.
- 23 Mokosuli, Y. S. (2008). Aktivitas antioksidasi dan antikanker ekstrak kulit batang Langsat (*Lansium domesticum* L.). (Tesis). Sekolah Pascasarjana Institut Pertanian Bogor.
- Mokosuli, Y. S. (2013). Karakter Morfologi, Sumber Pakan, Dan Bioaktivitas Farmakologis Racun Lebah Madu Endemik Sulawesi *Apis dorsata* Binghami DAN *Apis nigrocincta* Smith (HYMENOPTERA : APIDAE). (Disertasi). Program Pascasarjana Universitas Sam Ratulangi, Manado.
- Mokosuli, Y. S. & Repi, R. A. (2015). The Characteristics Of Bioactive Peptides And Antibacterial activity of honey bee (*Apis nigrocincta*) smith venom, endemic to sulawesi. *Molekul*, Vol. 10. No. 2. November: 135 - 144
- Mardawati, F., Filianty, & Marta, H. 2008. Kajian aktivitas antioksidan ekstrak kulit manggis (*Garcinia mangostana* L) dalam rangka pemanfaatan limbah kulit manggis di Kecamatan Puspahiang Kabupaten Tasikmalaya. Fakultas Teknologi Industri. Universitas Padjadjaran.
- 18 Nazzi, F. (2016). The hexagonal shape of the honeycomb cells depends on the construction behavior of bees. *Sci Rep*. 2016; 6: 28341. Published online 2016 Jun 20. doi: 10.1038/srep28341.
- 25 Otis, G.W. (1991). Revised distribution of three recently recognized species of honey bees in Asia. *Honeybee Science* 15:167–170.
- 60 Pasupuleti, V. R., Sammam, L., Ramesh, N., & Gan, S. H. (2017). Honey, Propolis, and Royal Jelly: A Comprehensive Review of Their Biological Actions and Health Benefits. *Oxidative Medicine and Cellular Longevity*. Volume 2017, Article ID 1259510, 21 pages <https://doi.org/10.1155/2017/1259510>
- Pérez-Pérez, E. M., Suárez, E., Peña-Vera, M. J., González, A. C., & Vit, P. (2013). Antioxidant activity and microorganisms in nest products of *Tetragonisca angustula* Latreille, 1811 from Mérida, Venezuela. pp. 1-8. In Vit P & Roubik DW, eds. *Stingless bees process honey and pollen in cerumen pots*. Facultad de Farmacia y Bioanálisis, Universidad de Los Andes; Mérida, Venezuela.
- 30 Raffiudin, R. (2002). Honey bee behavioural evolution and itpr gene structure studies. [PhD Thesis]. James Cook University <http://eprints.jcu.edu.au/1249>
- 4 Redha, A. (2010). Flavonoid: Struktur, Sifat Antioksidatif dan Peranannya Dalam Sistem Biologis. *Jurnal Berlin*, 9(2), 196–202. <https://doi.org/10.1186/2110-5820-1-7>
- Ra'ed, A., Ibrahim, K. N., Rula, M. D., & Mosa, A. (2008). Honey Bee Hive Modification for Propolis Collection. *Jordan Journal of Agricultural Sciences*, Vol 4(2)
- Repi, R. A., Mokosuli, Y.S., Ngangi, J., & Sumampouw, H. M. (2013). Hepatoprotective Activity Combination Between Morinda Citrifolia Linn (Mengkudu) Extract And Virgin Coconut Oil (VCO). *Journal of Biology, Agriculture and Healthcare*. Vol.3, No.13, pp 160-166.
- Rosli, N. L., Roslan, H., Oma, E. A., Mokhtar, N., Pit, N. H. A., & Asem N. (2016). Phytochemical analysis and antioxidant activities of *Trigona apicalis* propolis extract. *AIP Conference Proceedings* 1791, 020018 2016; doi: 10.1063/1.4968873. <https://aip.scitation.org/toc/apc/1791/1?expanded=1791>
- Sannora, V. B., Fonseca, M. A. D., & Silva L. (2005). Flavonoids and antiulcerogenic activity from *Byrsonima crassa* leaves extracts. *Journal of Ethnopharmacology*, vol.97, no.1,pp. 1–6,.
- Sawaya, A. C. H. F., Calado, J. C. P., Santos, L. C. D. S., Marcucci, M. C., Akatsu, I. P., & Soares, A. E. S., Abdelnur, V., Cunha I. B. D. S., and Eberlin M. N. (2009). Composition and antioxidant activity of propolis from three species of *Scaptotrigona* stingless bees. *Journal of ApiProduct and ApiMedical Science* 1 (2):

- 37 - 42 (2009). DOI 10.3896/IBRA.4.01.2.03
- Saavedra, N., Barrientos, L., Herrera, C. L., Alvear, M., Montenegro G., & Salazar L. A. (2011). Effect of Chilean propolis on cariogenic bacteria *Lactobacillus fermentum*. *Ciencia Investigacion Agraria*. [52]ine]. vol.38, n.1, pp.117-125. ISSN 0718-1620. <http://dx.doi.org/10.4067/S0718-16202011000100011>.
- Shalati [31] E. A., & Sanaa, M. M. S., (2012). Comparison of DPPH and ABTS assays for determining antioxidant potential of water and methanol extracts of *Spirulina platensis*. *Indian Journal of Geo-Marine Sciences* 42(5):556-564.
- Seal, T. (2016). Quantitative HPLC analysis of phenolic acids, flavonoids and ascorbic acid in four different solvent extracts of two wild edible leaves, *Sonchus arvensis* and *Oenanthe* [44]aris of North-Eastern region in India. *Journal of Applied Pharmaceutical Science*, 6(2), 157-166. <https://doi.org/10.7324/JAPS.2016.60225>.
- [37] Suseno, D., (2009). Aktivitas Antibakteri Propolis *Trigona* spp. Pada Dua Konsentrasi Berbeda [59]hadap Cairan Rumen Sapi. <https://repository.ipb.ac.id/bitstream/handle/123456789/12572/G09dsu1.pdfsequence=2>.
- Trusheva, B., Bankova, V., & Trunkova, D., (2007). Different extraction methods of biologically active components from propolis: A preliminary study. *Chemistry Central Journal*, 1:13 doi:10.1186/1752-153X-1-13.
- Veloz, J. J., Saavedra, N., Alvear, M., Zambrano [19] T., Barrientos, L., & Salazar, L. A. (2016). Polyphenol-Rich Extract from Propolis Reduces the Expression and Activity of Streptococcus mutans Glucosyltransferases at Subinhibitory [51]ncentration. *BioMed Research International* Volume 2016, Article ID 4302706, 7 pages <http://dx.doi.org/10.1155/2016/4302706>
- Waji, R. A., & Sugrani, A. (2009). Makalah Kimia Organik Bahan Alam Flavonoid (Quertecin). *Fakultas Matematika Dan Ilmu Pengetahuan Alam Universitas Hasanuddin*, 23.
- [15] Yang, D., Wang Q., Ke, L., Jiang J., & Ying T. (2007). antioxidant activities of various extracts of lotus [56]Nelumbo nucifera Gaertn) rhizome. *Asia Pacific Journal of Clinical Nutrition*. 16 (Suppl 1): 158-163.

The bioactive contents and antioxidant activity of honey bee nest extract of *Apis dorsata* Binghami from the North Sulawesi

ORIGINALITY REPORT

19%

SIMILARITY INDEX

13%

INTERNET SOURCES

11%

PUBLICATIONS

10%

STUDENT PAPERS

PRIMARY SOURCES

- 1** harvest.usask.ca 1%

Internet Source
- 2** Kandemir, İrfan, Mohammad G. Moradi, Berna Özden, and Ayça Özkan. "Wing geometry as a tool for studying the population structure of dwarf honey bees (*Apis florea* Fabricius 1876) in Iran", *Journal of Apicultural Research*, 2009. <1%

Publication
- 3** Maria L. Absy, André R. Rech, Marcos G. Ferreira. "Chapter 3 Pollen Collected by Stingless Bees: A Contribution to Understanding Amazonian Biodiversity", Springer Nature, 2018 <1%

Publication
- 4** Submitted to St. Ursula Academy High School <1%

Student Paper
- 5** Submitted to Georgia State University <1%

Student Paper

6	athmsi.org Internet Source	<1 %
7	Submitted to Karadeniz Teknik University Student Paper	<1 %
8	Submitted to University Tun Hussein Onn Malaysia Student Paper	<1 %
9	ejournal.unib.ac.id Internet Source	<1 %
10	etd.lib.ttu.edu Internet Source	<1 %
11	jurnal.yudharta.ac.id Internet Source	<1 %
12	Submitted to Politeknik STIA LAN Student Paper	<1 %
13	Submitted to Universitas Sultan Ageng Tirtayasa Student Paper	<1 %
14	Submitted to Australian National University Student Paper	<1 %
15	www.notulaebiologicae.ro Internet Source	<1 %
16	Submitted to The Robert Gordon University Student Paper	<1 %

- | | | |
|----|--|------|
| 17 | www.eolss.net
Internet Source | <1 % |
| 18 | Cheng-Kai Liu, Chi-Tang Huang, Jyun-Cong Lai, Ko-Ting Cheng. "Spontaneous and field-induced evolutions of 2D patterns in fingerprint chiral textures", <i>Journal of Molecular Liquids</i> , 2022
Publication | <1 % |
| 19 | Mariya Gover Antoniraj, Kasi Pandima Devi, Ioana Berindan-Neagoe, Seyed Fazel Nabavi et al. "Oral microbiota in cancer: could the bad guy turn good with application of polyphenols?", <i>Expert Reviews in Molecular Medicine</i> , 2022
Publication | <1 % |
| 20 | ir.jkuat.ac.ke
Internet Source | <1 % |
| 21 | sciencescholar.us
Internet Source | <1 % |
| 22 | www.smujo.id
Internet Source | <1 % |
| 23 | Josepin P Konda, Jainer P Siampa, Trina E Tallei, Billy J Kepel, Fatimawali Fatimawali. "Aktivitas Antioksidan Ekstrak Metanol Biji Langsung (<i>Lansium domesticum</i> var. <i>pubescens</i>) dan Duku (<i>Lansium domesticum</i>) | <1 % |

var. domesticum) dengan Metode DPPH",

JURNAL ILMIAH SAINS, 2020

Publication

24

jurnalnasional.ump.ac.id

Internet Source

<1 %

25

www.bio.usyd.edu.au

Internet Source

<1 %

26

Submitted to University of Alabama,
Huntsville

Student Paper

<1 %

27

www.msmbb.org.my

Internet Source

<1 %

28

Submitted to Napier University

Student Paper

<1 %

29

Rolef Rumondor, Muhammad Rino Komalig,
Kamaluddin Kamaluddin. "Efek Pemberian
Ekstrak Etanol Daun Leilem (Clerodendrum
minahasae) terhadap Kadar Kreatinin, Asam
Urat dan Ureum pada Tikus Putih (Rattus
novergicus)", Bio-Edu: Jurnal Pendidikan
Biologi, 2019

Publication

<1 %

30

journal.biotrop.org

Internet Source

<1 %

31

jurnal.ugm.ac.id

Internet Source

<1 %

32	repository.poltekkespim.ac.id Internet Source	<1 %
33	www.astesj.com Internet Source	<1 %
34	Alfin Surya, Yulia Yesti. "AKTIVITAS ANTIOKSIDAN EKSTRAK METANOL KULIT JENGKOL (Pithecellobium jiringa) DENGAN TIGA WAKTU MASERASI", Human Care Journal, 2018 Publication	<1 %
35	Submitted to Higher Education Commission Pakistan Student Paper	<1 %
36	ediss.uni-goettingen.de Internet Source	<1 %
37	issuu.com Internet Source	<1 %
38	Submitted to Universitas Jenderal Soedirman Student Paper	<1 %
39	balimedicaljournal.org Internet Source	<1 %
40	edoc.pub Internet Source	<1 %
41	hero.epa.gov Internet Source	<1 %

- | | | |
|----|---|------|
| 42 | chemicalbulletin.upt.ro
Internet Source | <1 % |
| 43 | Awunyo-Vitor, Dadson, Shaibu Ishak, and Godfred Seidu Jasaw. "Urban Households' Willingness to Pay for Improved Solid Waste Disposal Services in Kumasi Metropolis, Ghana", Urban Studies Research, 2013.
Publication | <1 % |
| 44 | Submitted to Stockholm University
Student Paper | <1 % |
| 45 | Venegas, Yovani, Carlos Peña, Edgar Pastene, and David Contreras. "A new near-infrared method for simultaneous determination of caffeic acid phenethyl ester and antioxidant activity of propolis samples", Journal of Apicultural Research, 2016.
Publication | <1 % |
| 46 | journal.unhas.ac.id
Internet Source | <1 % |
| 47 | www.amdi.usm.my
Internet Source | <1 % |
| 48 | www.imedpub.com
Internet Source | <1 % |
| 49 | Kharis Sugiarto, Muhammad Rivai, Astria Nur Irfansyah. "Control of Livestock Waste Odors Using Gas Sensors and Fuzzy Logic", 2019 | <1 % |

12th International Conference on Information & Communication Technology and System (ICTS), 2019

Publication

50

Submitted to The Institute for Optimum Nutrition

Student Paper

<1 %

51

Submitted to University of Glamorgan

Student Paper

<1 %

52

lookformedical.com

Internet Source

<1 %

53

qiup.edu.my

Internet Source

<1 %

54

Katarzyna Paradowska, Agnieszka Zielińska, Marzena Kuras, Iwona Wawer. " The composition of bee pollen color fractions evaluated by solid-state H and C NMR: their macroelement content and antioxidant properties ", Journal of Apicultural Research, 2017

Publication

<1 %

55

foodsofthecape.blogspot.com

Internet Source

<1 %

56

koreascience.or.kr

Internet Source

<1 %

57	Kohji Takara. "Effects of propolis extract on sensitivity to chemotherapeutic agents in HeLa and resistant sublines", <i>Phytotherapy Research</i> , 09/2007 Publication	<1 %
58	repository.stifarm-padang.ac.id Internet Source	<1 %
59	www.atlantis-press.com Internet Source	<1 %
60	Submitted to University of Ulster Student Paper	<1 %
61	fmipa.umri.ac.id Internet Source	<1 %
62	guaiaca.ufpel.edu.br:8080 Internet Source	<1 %
63	jos.unsoed.ac.id Internet Source	<1 %
64	repository.uhamka.ac.id Internet Source	<1 %
65	repository.unisba.ac.id Internet Source	<1 %
66	www.pjps.pk Internet Source	<1 %

67

F.E Hasibuan, W Syahfitri, S Ilyas, S Hutahaean. " Phytochemical screening, antioxidant activity and thin-layer chromatography test of methanol extract and simplicia leaves of loquat (Lindl) ", IOP Conference Series: Materials Science and Engineering, 2020

Publication

<1 %

68

Ivan Gunawan, Iwan Vanany, Erwin Widodo. "Cost-benefit model in improving traceability system: case study in Indonesian bulk-liquid industry", Supply Chain Forum: An International Journal, 2019

Publication

<1 %

69

Park, Yong-Seo, Korsak Towantakavanit, Jacek Namiesnik, Magdalena Sliwka-Kaszynska, Barbara Kuszniereicz, Katarzyna Kozlowska-Tylingo, Hanna Leontowicz, Maria Leontowicz, Alma Leticia Matrinez-Ayala, and Shela Gorinstein. "Partial Characterization of Three Korean White Lotus Cultivars", Journal of Agricultural and Food Chemistry, 2009.

Publication

<1 %

Exclude quotes Off

Exclude matches < 2 words

Exclude bibliography Off